

## Review

### Correspondence

D. Robertson

d.robertson@dental.gla.ac.uk

# The microbiology of the acute dental abscess

D. Robertson<sup>1</sup> and A. J. Smith<sup>2</sup>

<sup>1</sup>Department of Restorative Dentistry, University of Glasgow, Medical Faculty, Glasgow Dental Hospital and School, 378 Sauchiehall Street, Glasgow G2 3JZ, UK

<sup>2</sup>Infection and Immunity Section, University of Glasgow, Medical Faculty, Glasgow Dental Hospital and School, 378 Sauchiehall Street, Glasgow G2 3JZ, UK

The acute dental abscess is frequently underestimated in terms of its morbidity and mortality. The risk of potential serious consequences arising from the spread of a dental abscess is still relevant today with many hospital admissions for dental sepsis. The acute dental abscess is usually polymicrobial comprising facultative anaerobes, such as viridans group streptococci and the *Streptococcus anginosus* group, with predominantly strict anaerobes, such as anaerobic cocci, *Prevotella* and *Fusobacterium* species. The use of non-culture techniques has expanded our insight into the microbial diversity of the causative agents, identifying such organisms as *Treponema* species and anaerobic Gram-positive rods such as *Bulleidia extructa*, *Cryptobacterium curtum* and *Mogibacterium timidum*. Despite some reports of increasing antimicrobial resistance in isolates from acute dental infection, the vast majority of localized dental abscesses respond to surgical treatment, with antimicrobials limited to spreading and severe infections. The microbiology and treatment of the acute localized abscess and severe spreading odontogenic infections are reviewed.

## Introduction

The acute dental abscess is frequently underestimated in terms of its morbidity and mortality. The acute dental abscess usually occurs secondary to dental caries, trauma or failed root treatment. After the intact pulp chamber is breached, colonization of the root canals occurs with a diverse mix of anaerobic bacteria. The walls of the necrotic root canals become colonized by a specialized mixed anaerobic biofilm (Chavez de Paz, 2007). While asymptomatic necrosis is common, abscess formation occurs when these bacteria and their toxic products enter the periapical tissues via the apical foramen and induce acute inflammation and pus formation (Nair, 2004). The root canal microbiota is the main stimulus for the development of acute symptoms. The main signs and symptoms of the acute dental abscess (often referred to as a periapical abscess or infection) are pain, swelling, erythema and suppuration usually localized to the affected tooth, although the abscess can frequently spread causing a spreading odontogenic infection which can be accompanied by sepsis syndrome. The role of bacteria in the pathogenesis of the lesion is undisputed but modern diagnostic techniques have not identified a single causative pathogen. The dentoalveolar abscess is polymicrobial comprising various facultative anaerobes, such as the viridans group streptococci and the *Streptococcus anginosus* group, and strict anaerobes, especially anaerobic cocci,

*Prevotella* and *Fusobacterium* species. The presence of anaerobes both cultivable and uncultivable tends to predominate. The vast majority of dental abscesses respond to surgical treatment, such as drainage of pus and elimination of the source of infection, with antibiotic use limited to severe spreading infections. The microbiology of the acute dentoalveolar abscess and its treatment in the light of improved culture and diagnostic methods are reviewed.

## Background

Historically, the potential for a dental abscess to spread causing severe sepsis and death has been known since antiquity although the role of bacteria in this process was not recognized until the turn of the 20th century (Turner Thomas, 1908). When the Bills of Mortality (London) began listing the causes of death in the early 1600s, 'teeth' were listed as the 5th or 6th leading cause of death (Clarke, 1999). By the turn of the 20th century, dental infections were associated with a mortality rate of 10–40% (Turner Thomas, 1908). Collecting appropriate current epidemiological data is difficult due to differences in the methods of reporting throughout the world but the evidence suggests that it is still a significant clinical problem. In Scotland there were 3500 hospital admissions between 2000 and 2005 for acute dental infections while hospitals in England saw a doubling of admissions for surgical treatment of dental abscesses over a similar period

The authors contributed equally to this paper.

(Information & Statistics Division, 2007; Thomas *et al.*, 2008). In the United States, despite a generally dentally motivated population a large prospective study reported that 13 % of adult patients sought treatment for dental pain and infection over a 24 month follow-up (Boykin *et al.*, 2003), and in the US hospital admissions relating to acute dental infections are estimated to occur at rate of 1 per 2600 head of population per year (Wang *et al.*, 2005). Periapical abscesses account for 47 % of all dental-related attendances at paediatric emergency rooms in the United States (Graham *et al.*, 2000). While the studies cited above do not give an exhaustive analysis of the extent of this problem in different parts of the world, nonetheless they do lend credence to the view that dental infections have significant implications for patient morbidity and health care systems. Improved methods of reporting of this common infection should be developed in order to allow more detailed epidemiological analysis to be carried out.

### Cultural analysis of the acute dental abscess

Cultural analysis remains the backbone of clinical practice and the findings of a number of prospective and retrospective studies give a valuable insight into the bacteria which are often present. Efforts to identify the causative pathogens involved in the development of the dental abscess have in the past been hampered by inappropriate methods of sampling. The ideal clinical sample from an acute dental abscess is an aspirate through intact mucosa disinfected by an appropriate antiseptic mouthwash or swab, e.g. chlorhexidine, although some researchers have sampled purulent exudates from within infected canals (Lewis *et al.*, 1990; Chavez de Paz Villanueva, 2002). This will reduce contamination from the normal oral flora. Previous studies using swabs of purulent material have demonstrated poor recovery of strict anaerobes and low mean numbers of isolates per sample (range 1.0–1.6) (Lewis *et al.*, 1990). Pure cultures from an acute dental abscess are unusual (Reader *et al.*, 1994), and mixed aerobic infections are also uncommon, accounting for 6 % of abscesses (Goumas *et al.*, 1997). Dental abscesses caused solely by strict anaerobes occur in approximately 20 % of cases although there is a wide range depending on recovery conditions (6–63 %) (Brook *et al.*, 1991; Gorbach *et al.*, 1991; Goumas *et al.*, 1997; Khemaleelakul *et al.*, 2002).

A complex mix of strict anaerobes and facultative anaerobes accounts for most infections (59–75 %), which can prove challenging to non-specialist microbiology laboratories (Gorbach *et al.*, 1991; Goumas *et al.*, 1997; Kuriyama *et al.*, 2000a). In mixed infections, strict anaerobes outnumber facultatives by a ratio which varies between 1.5–3:1, again depending on the recovery and culture conditions (Baumgartner & Xia, 2003; Khemaleelakul *et al.*, 2002; Kulekci *et al.*, 1996; Lewis *et al.*, 1993; Roche & Yoshimori, 1997; Sakamoto *et al.*, 1998). The mean number of species recovered by culture from dentoalveolar

aspirates is 4 with a range of between 1 and 7.5 (Fazakerley *et al.*, 1993; Khemaleelakul *et al.*, 2002; Reader *et al.*, 1994).

### Facultative anaerobes

The most commonly found facultative anaerobes belong to the viridans group streptococci and the *anginosus* group streptococci. Difficulties lie in interpretation of the literature on the relative contributions of individual species from these two major groups of organisms due to taxonomic changes and accuracy of identification. The viridans group streptococci comprise the *mitis* group, *oralis* group, *salivarius* group, *sanguinis* group and the *mutans* group (Facklam, 2002). The *anginosus* group (formerly referred to as '*Streptococcus milleri*' or *Streptococcus anginosus*) is also identified and reported with varying degrees of accuracy ranging from just beta-haemolytic streptococci down to species level.

Historically, *Staphylococcus* species have not been considered members of the oral flora or to play a major role in the pathogenesis of oral infections. However, a number of more recent studies have indicated that staphylococci may indeed be a more frequent colonizer of the oral tissues than previously thought (Smith *et al.*, 2001). The recovery rates using conventional culture of *Staphylococcus aureus* from the acute dental abscess range from 0.7 to 15 % (Brook *et al.*, 1991; Goumas *et al.*, 1997; Kulekci *et al.*, 1996; Kuriyama *et al.*, 2002b; Roche & Yoshimori, 1997; Siqueira *et al.*, 2001d) although some workers have noted higher recovery rates of 47 % (Mangundjaja & Hardjwinata, 1990). Interestingly, *Staphylococcus aureus* has been reported to occur more frequently in severe dental abscesses from children (Brook *et al.*, 1991; Coticchia *et al.*, 2004; Coulthard & Isaacs, 1991; Dodson *et al.*, 1989; Tan *et al.*, 2001).

Recovery rates of coagulase-negative strains of staphylococci (usually reported as *Staphylococcus epidermidis*) are generally higher with figures ranging from 4 to 65 % (Gorbach *et al.*, 1991; Goumas *et al.*, 1997; Khemaleelakul *et al.*, 2002; Kuriyama *et al.*, 2002b; Lewis *et al.*, 1995; Mangundjaja & Hardjwinata, 1990; Sakamoto *et al.*, 1998; Storoe *et al.*, 2001). *Staphylococcus* species may also be associated with refractory infections not responding to endodontic treatment (Reader *et al.*, 1994).

### Anaerobes

Similar difficulties exist for cross-study comparisons of identification and prevalence of strict anaerobes. The most commonly isolated genera include anaerobic streptococci, *Fusobacterium* species and the black-pigmented anaerobes such as *Prevotella* and *Porphyromonas* species (Sundqvist *et al.*, 1989).

The nomenclature and recent changes in taxonomy have complicated the comparison of more recent studies with older studies due to the renaming of several species, specifically the *Prevotella*, *Bacteroides* and *Porphyromonas*

species. An important group of pathogens that has undergone much in the way of taxonomic rearrangement, this group, often referred to as the 'oral *Bacteroides*' and black-pigmenting anaerobes group, has been reclassified. The *Bacteroides* species have been divided into the saccharolytic genus *Prevotella* and the asaccharolytic genus *Porphyromonas*. The genus *Bacteroides* has been restricted to the fermentative *Bacteroides fragilis* and its closely related species. *B. fragilis*, a more common isolate from intra-abdominal infections, has only infrequently been reported from acute dentoalveolar infections and is not regarded as an oral commensal. The member of the *Bacteroides* genus most likely to be recovered from an acute dental abscess is *Bacteroides forsythus* (now transferred to a new genus as *Tannerella forsythia*) (Gomes *et al.*, 2006).

The most commonly reported anaerobic Gram-negative bacilli from acute dentoalveolar infections are species from the pigmented *Prevotella intermedia* group (comprising *Prevotella intermedia*, *Prevotella nigrescens* and *Prevotella pallens*), *Porphyromonas endodontalis* and *Porphyromonas gingivalis* (Jacinto *et al.*, 2006). The *Prevotella* species are the most frequent isolates, found in 10–87% of dentoalveolar abscesses (Baumgartner *et al.*, 2004; Fazakerley *et al.*, 1993; Kolokotronis, 1999; Kulekci *et al.*, 1996; Kuriyama *et al.*, 2005; Lewis *et al.*, 1993; Riggio *et al.*, 2006; Roche & Yoshimori, 1997; Sakamoto *et al.*, 1998; Siqueira *et al.*, 2001b, d; Wade *et al.*, 1994).

The genus *Fusobacterium* is frequently reported in infections of the head and neck with reports indicating that *Fusobacterium* species can be detected in up to 52% of specimens (Gill & Scully, 1990; Gilmore *et al.*, 1988; Gorbach *et al.*, 1991; Goumas *et al.*, 1997; Kulekci *et al.*, 1996; Kuriyama *et al.*, 2000a, b, 2005, 2006; Lewis *et al.*, 1993; Mangundjaja & Hardjwinata, 1990; Sakamoto *et al.*, 1998; Wade *et al.*, 1994). Taxonomy and nomenclature of the genus *Fusobacterium* also cause difficulties in comparisons across studies. Within the human oral flora, *Fusobacterium periodonticum* and *Fusobacterium nucleatum* (which includes subsp. *nucleatum*, subsp. *polymorphum*, subsp. *animalis*, subsp. *vincentii* and subsp. *fusiforme*) are frequently detected with *F. nucleatum* recovered most frequently from the acute dental abscess (Dzink *et al.*, 1990; Chavez de Paz Villanueva, 2002; Sassone *et al.*, 2008). Studies utilizing non-culture techniques for analysis of the dental abscess for the presence of *F. nucleatum* have reported a prevalence of 73% (Baumgartner *et al.*, 2004).

The Clostridia are infrequently reported from odontogenic infections either as a sole pathogen or as part of the abscess flora. Workers have recovered *Clostridium* species from 2–20% of specimens (Gorbach *et al.*, 1991; Goumas *et al.*, 1997; Khemalelakul *et al.*, 2002; Roche & Yoshimori, 1997). Where speciated, these isolates have included *Clostridium hastiforme*, *Clostridium histolyticum*, *Clostridium perfringens*, *Clostridium subterminale* and *Clostridium clostridioforme* (Khemalelakul *et al.*, 2002; Roche & Yoshimori, 1997). Although other *Clostridium*

species such as *Clostridium sporogenes*, *Clostridium bifermentans*, *Clostridium botulinum*, '*Clostridium oedomatiens*' and '*Clostridium welchii*' have been recovered from carious dentine, they appear to be infrequent pathogens in the oral cavity (Van Reenan & Coogan, 1970).

### Analysis of the microflora of the acute dental abscess using molecular biological techniques

Close attention to specimen collection and processing on selective and non-selective agars under appropriate atmospheric conditions has improved the routine diagnostic yield from acute dental abscesses. However, despite meticulous attention to detail, it is apparent that many genera of bacteria have yet to be cultured from many infectious diseases including the acute dental abscess (Siqueira & Rocas, 2005). The use of culture-independent or molecular diagnostic techniques has expanded our insight into the microbial ecology of the dental abscess. There is an increasing reliance on genetic methods of identification with 16S rRNA gene sequencing frequently being used for research purposes. Broadly speaking, the molecular analysis may take one of two approaches. Firstly, the use of molecular cloning and sequencing techniques to identify uncultivable micro-organisms using 16s rRNA or rDNA has led to the identification of several novel species (Dymock *et al.*, 1996). The second approach utilizes PCR or DNA–DNA hybridization checkerboard techniques (Siqueira *et al.*, 2001d, 2002a) and more recently 16S rRNA gene sequencing and species-specific primers searching for the presence of specific microbes (Dymock *et al.*, 1996; Riggio *et al.*, 2006; Rocas & Siqueira, 2005; Sakamoto *et al.*, 2006; Siqueira *et al.*, 2001b, c, 2002b, 2003). This approach has demonstrated the higher prevalence of more fastidious organisms such as *Treponema* species in the acute dental abscess.

*Treponema* species are strictly anaerobic, motile, helically shaped bacteria. Within the oral cavity they are more usually associated with diseases of the periodontium. There are a number of different species described from the oral cavity including *Treponema amylovorum*, *Treponema denticola*, *Treponema maltophilum*, *Treponema medium*, *Treponema pectinovorum*, *Treponema socranskii* and '*Treponema vincentii*' (Chan & McLaughlin, 2000). The treponemes are difficult to cultivate and differentiate and only *T. denticola*, *T. pectinovorum*, *T. socranskii* and '*T. vincentii*' have been readily cultivated. Recent work using PCR detection has indicated a surprisingly high prevalence of *Treponema* species within the acute dental abscess. Siqueira & Rocas (2004c) found that *T. denticola* was present in up to 79% of dental abscesses, with lower detection rates reported by other workers (Baumgartner *et al.*, 2003; Siqueira *et al.*, 2001a, c; Gomes *et al.*, 2006; Cavrini *et al.*, 2008). Other *Treponema* species were found in lower numbers, including *T. socranskii* (in 26% of aspirates), *T. pectinovorum* (14–21% of aspirates), *T. amylovorum* (16% of aspirates) and *T. medium* (5% of

aspirates). Other species such as *Treponema lecithinolyticum*, '*T. vincentii*' and *T. maltophilum* were not detected. Despite the possession of a number of potential virulence factors, the precise role of these poorly understood and under-reported organisms in the pathogenesis of the acute dental abscess is unclear.

### Unfamiliar species

Improvements in sampling, culture and identification have led to a greater insight into the diversity of the microbial flora in an acute dental abscess. This has resulted in the reporting of micro-organisms which are probably more accurately described as 'unfamiliar' rather than 'new' implying their recent appearance. These include members of the genus *Atopobium* (Gram-positive strictly anaerobic coccobacilli), for example *Atopobium parvulum* and *Atopobium rimae*. Anaerobic Gram-positive rods include *Bulleidia extracta*, *Cryptobacterium curtum*, *Eubacterium sulci*, *Mogibacterium timidum* and *Mogibacterium vesicum* (Sakamoto *et al.*, 2006), *Pseudoramibacter alactolyticus* and *Slakia exigua* (Siqueira & Rocas, 2003c).

Other unfamiliar species include anaerobic Gram-negative rods such as *Filifactor alocis* (Siqueira & Rocas, 2003a, 2004b; Gomes *et al.*, 2006) and *Dialister pneumosintes* (Siqueira *et al.*, 2005; Siqueira & Rocas, 2003b, 2004b). *Centipeda periodontii* and *Selenomonas sputigena* are multi-flagellated, motile, anaerobic, Gram-negative rods also found recently in the acute dental abscess (Siqueira & Rocas, 2004a). *Catonella morbi*, a Gram-negative anaerobe formerly known as *Bacteroides* D42, was found in 16% of 19 aspirates, and *Granulicatella adiacens*, a facultative anaerobic Gram-positive coccus formerly known as nutritionally variant streptococci, was present in 11% of 19 aspirates (Rocas & Siqueira, 2005; Siqueira & Rocas, 2006).

The detection of these unfamiliar species has opened up a whole new area for possible study into the virulence factors possessed by these bacteria and their relative influence on the pathogenesis of the acute dental abscess and interactions with more commonly isolated and better understood pathogens. These techniques are not without their limitations and meticulous asepsis is required throughout the sampling and analysis procedure to avoid contamination due to the sensitivity of these methods. Furthermore, until recently these techniques could only give semiquantitative analysis of aspirates and indeed some papers cited above can only show the presence or absence of the species in question. This will improve with the advent of quantitative real-time PCR. The use of species-specific primers targeting the 16S rRNA gene or similar is also limited by the fact that they cannot distinguish between transcriptionally active viable cells and those non-vital bystanders. Advanced molecular techniques using reverse transcriptase are finding methods of overcoming these limitations currently. Finally, molecular techniques can give little useful information to guide the clinician in the choice of antibiotic if required.

### Antibiotic resistance

The antibiotics most commonly prescribed in the management of acute dental abscesses in the UK are amoxicillin, penicillin, metronidazole and erythromycin (Palmer *et al.*, 2000). Antibiotic resistance in microbes recovered from the acute dental abscess has been reported to be increasing (with the exception of metronidazole) in some populations studied over the last few decades (Kuriyama *et al.*, 2006; Lewis *et al.*, 1989, 1995; Storie *et al.*, 2001). Care must be taken in interpretation of studies due to differences in details of the identification of isolates, selection of appropriate breakpoints and their relevance to oral infections and lack of details on MICs (for example, not all studies report MIC<sub>90</sub> data). This is illustrated by reports of resistance rates for amoxicillin ranging from 9 to 54% of common isolates from acute dental abscesses (Baumgartner & Xia, 2003; Gilmore *et al.*, 1988; Khemalelakul *et al.*, 2002; Kuriyama *et al.*, 2000a, 2002a, 2005; Lewis *et al.*, 1993, 1995; Smith & Jackson, 2003).

However, analysis of a number of reports does reveal a trend that the least susceptible isolates from an acute abscess are more likely to be black-pigmented *Prevotella* species, such as *Prevotella intermedia*, *Prevotella melaninogenica*, *Prevotella denticola* and *Prevotella loescheii*, followed by the non-pigmenting *Prevotella* species, such as *Prevotella oralis*, *Prevotella buccae*, *Prevotella disiens* and *Prevotella bivia*. The most common mechanism of antimicrobial resistance detected is production of beta-lactamase, probably related to the group 2e class of beta-lactamases (Valle *et al.*, 1998). This group of beta-lactamases demonstrates a major activity on cephalosporins rather than penicillins and retains susceptibility to inhibition by clavulanic acid and tazobactam. The presence of beta-lactamase production in other anaerobes from acute dental infections, such as *Porphyromonas* and *Fusobacterium* species, appears infrequent.

Studies have reported that resistance from the *anginosus* group of streptococci is rare with 1 of 43 isolates (2.3%) having a penicillin MIC greater than 1 mg l<sup>-1</sup> and 7 of 45 (15.5%) 'viridans group' streptococci having a penicillin MIC greater than 1 mg l<sup>-1</sup> (Lewis *et al.*, 1995). More recently, this group has failed to detect penicillin resistance by NCCLS breakpoints in 64 streptococcal isolates (Kuriyama *et al.*, 2005). Reduced susceptibility to penicillin is more prevalent in the *mitis* group streptococci than in the *anginosus* group (Smith & Jackson, 2003).

Resistance to macrolides appears to have a higher prevalence in the 'viridans group streptococci', anaerobic streptococci and *Prevotella* species (Kuriyama *et al.*, 2000a, 2001, 2002b). This is linked to penicillin resistance. The erythromycin MIC<sub>90</sub> of 139 streptococcal isolates was 1.0 mg l<sup>-1</sup> for penicillin-susceptible isolates and 2.0 mg l<sup>-1</sup> for penicillin-resistant isolates (Kuriyama *et al.*, 2000a). For pigmented *Prevotella* species (*n*=93), the erythromycin MIC<sub>90</sub> for penicillin-susceptible isolates was 8 mg l<sup>-1</sup> and for penicillin non-susceptible isolates was 64 mg l<sup>-1</sup> (Kuriyama *et al.*,

2000a). Both penicillin-susceptible and -resistant isolates of *Fusobacterium* species ( $n=90$ ) had an MIC<sub>90</sub> of 64 mg l<sup>-1</sup> for erythromycin. Macrolide resistance is most commonly due to acquisition of one of a number of *erm* genes (erythromycin methylases resulting in reduced binding of macrolides to the 50S ribosomal subunit).

The prevalence of resistance to lincosamides, such as clindamycin, is low. For example, Kuriyama *et al.* (2000a) reported on 664 isolates from 163 patients and found a clindamycin MIC<sub>90</sub> of 2 mg l<sup>-1</sup> for 15 strains of penicillin-resistant anaerobic streptococci, with all the remaining isolates (649) having a clindamycin MIC<sub>90</sub> less than 0.5 mg l<sup>-1</sup>. Similar low levels of clindamycin resistance have been found in the UK (Kuriyama *et al.*, 2005) and elsewhere (Kuriyama *et al.*, 2001).

### Severe odontogenic infection

Deep neck and mediastinal abscesses are a rare complication of the dental abscess but spread of odontogenic infections accounts for up to 57% of deep neck abscesses (Mihos *et al.*, 2004; Sancho *et al.*, 1999). With the potential for infection spreading to the interpleural space and mediastinal tissue, the mortality rate of mediastinitis continues to be 17–50% despite aggressive use of antibiotics and advances in intensive care facilities (Corsten *et al.*, 1997; Marty-Ane *et al.*, 1999). Death usually occurs due to sepsis and multi-organ failure although airway occlusion is also a significant complication and requires early management by tracheostomy. Host factors affected by the patient's general health condition play a significant role. Specific at-risk groups are diabetics (Jimenez *et al.*, 2004; Tung-Yiu *et al.*, 2000) and the elderly (Wang *et al.*, 2003). Most studies report that males are affected more often by severe odontogenic infections than females in both adult (Flynn *et al.*, 2006; Sancho *et al.*, 1999) and paediatric (Dodson *et al.*, 1989) groups.

The bacteriology of the spreading odontogenic infection appears subtly different from that of the more localized dental abscess with *anginosus* group streptococci and *Fusobacterium* species forming the predominant flora isolated (Han & Kerschner, 2001; Heimdahl *et al.*, 1985; Schuman & Turner, 1999). Recent work suggests a significant role for *Prevotella* species in the spreading odontogenic infection with these species accounting for 50% of clones analysed from samples of pus aspirated from spreading odontogenic infections (Riggio *et al.*, 2006). Deep neck infections in the paediatric population also appear to be significantly different bacteriologically, with an increased involvement of *Staphylococcus aureus* (Brook, 1987; Coticchia *et al.*, 2004; Coulthard & Isaacs, 1991; Dodson *et al.*, 1989; Tan *et al.*, 2001) and group A streptococci (Coticchia *et al.*, 2004).

### Treatment of the acute dental abscess

There is no consensus over gold standard treatment as evidenced by the wide variety in endodontic and surgical

protocols and antibiotic prescribing (Kuriyama *et al.*, 2005). This ongoing controversy is a result of a lack of sufficient evidence to support the use of one antibiotic regimen over another or to indicate one treatment modality over another. Clinical trials in the treatment of the dental abscess are often flawed in design, limiting validity and applicability of results. Many studies are inadequately blinded and have not adequately controlled inclusion criteria. This has resulted in patients receiving multiple surgical and medical interventions making it impossible to analyse the relative contributions of each intervention to the success of treatment (Adriaenssen, 1998; Davis & Balcom, 1969; Fazakerley *et al.*, 1993; Fouad *et al.*, 1996; Gilmore *et al.*, 1988; Hanna, 1991; Kuriyama *et al.*, 2005; Lewis *et al.*, 1986, 1993; Mangundjaja & Hardjwinata, 1990). In one study, a total of six antibiotic regimens were prescribed with patients also receiving either incision and drainage or drainage through the tooth (Kuriyama *et al.*, 2005). Other difficulties include a lack of standardization in how the primary outcome is measured, with some studies relying on clinical assessment of relatively crude measures, i.e. failure, slight improvement, cure (Adriaenssen, 1998; Gilmore *et al.*, 1988; Hanna, 1991; Kuriyama *et al.*, 2005), and others a combination of both patient responses and clinical examination with well-defined but subjective criteria (Fazakerley *et al.*, 1993; Fouad *et al.*, 1996; Mangundjaja & Hardjwinata, 1990; Paterson & Curzon, 1993; Schuen *et al.*, 1974; von Konow *et al.*, 1992).

The above criticisms highlight the need for good-quality clinical trials to be held of sufficient size and scientific rigor to answer the question about the ideal treatment of the acute dental abscess. Nonetheless there are a number of recommendations which can be suggested based on the current evidence. Antibiotics should only be prescribed in patients exhibiting signs of local spread or systemic involvement. There is currently insufficient evidence to advocate the use of one regimen over another; however, a high dose used for as short a course as is consistent with a clinical cure has been shown to be effective and may reduce the development of resistance (Lewis *et al.*, 1986). If empirical antibiotics are required, the following may be considered. Amoxicillin remains the antimicrobial of first choice. If local patterns of antimicrobial resistance indicate a high prevalence of resistance to amoxicillin then the use of either metronidazole (Roche & Yoshimori, 1997) or amoxicillin in combination with clavulanic acid (Lewis *et al.*, 1993) should be considered as alternatives. Clindamycin remains an alternative in individuals who are allergic to the penicillin group of antibiotics (Gilmore *et al.*, 1988; Mangundjaja & Hardjwinata, 1990; Schuen *et al.*, 1974).

### Conclusions

With the advent of molecular diagnostic techniques, our insight into the diversity of the polymicrobial collection that comprises the dental abscess is expanding. Factors

influencing the process of bacterial succession from the saccharolytic cariogenic flora to the more anaerobic and proteolytic flora of the dental abscess remain unknown. Determination of factors that influence the spread of infection from a localized collection at the apex of a tooth to a cellulitis and life-threatening sepsis would aid treatment decisions. There are surprisingly few well-controlled studies into the most appropriate treatment regimen for the acute dental abscess, with most of the evidence pointing towards a key role for prompt surgical intervention and timely review. Antimicrobials should be reserved for patients with evidence of cellulitis and signs of sepsis. Although comparing different studies on antimicrobial susceptibility from dental abscesses is difficult, available data suggest that at present most isolates are still susceptible to first-line beta-lactam agents. In current times where the overuse of the prescription pad has eroded the antimicrobial arsenal, more effort could be placed on appropriate antimicrobial prescribing for the treatment of acute dental infections. Alternatively, strategies to improve oral health and reduce the incidence of dental caries, the main cause of the dental abscess, would maximize use of resources.

## References

- Adriaenssen, C. F. (1998).** Comparison of the efficacy, safety and tolerability of azithromycin and co-amoxiclav in the treatment of acute periapical abscesses. *J Int Med Res* **26**, 257–265.
- Baumgartner, J. C. & Xia, T. (2003).** Antibiotic susceptibility of bacteria associated with endodontic abscesses. *J Endod* **29**, 44–47.
- Baumgartner, J. C., Khemaleekul, S. U. & Xia, T. (2003).** Identification of spirochetes (treponemes) in endodontic infections. *J Endod* **29**, 794–797.
- Baumgartner, J. C., Siqueira, J. F., Jr, Xia, T. & Rocas, I. N. (2004).** Geographical differences in bacteria detected in endodontic infections using polymerase chain reaction. *J Endod* **30**, 141–144.
- Boykin, M. J., Gilbert, G. H., Tilashalski, K. R. & Shelton, B. J. (2003).** Incidence of endodontic treatment: a 48-month prospective study. *J Endod* **29**, 806–809.
- Brook, I. (1987).** Microbiology of retropharyngeal abscesses in children. *Am J Dis Child* **141**, 202–204.
- Brook, I., Frazier, E. H. & Gher, M. E. (1991).** Aerobic and anaerobic microbiology of periapical abscess. *Oral Microbiol Immunol* **6**, 123–125.
- Cavrini, F., Pirani, C., Foschi, F., Montebugnoli, L., Sambri, V. & Prati, C. (2008).** Detection of *Treponema denticola* in root canal systems in primary and secondary endodontic infections. A correlation with clinical symptoms. *New Microbiol* **31**, 67–73.
- Chan, E. C. & McLaughlin, R. (2000).** Taxonomy and virulence of oral spirochetes. *Oral Microbiol Immunol* **15**, 1–9.
- Chavez De Paz, L. E. (2007).** Redefining the persistent infection in root canals: possible role of biofilm communities. *J Endod* **33**, 652–662.
- Chavez de Paz Villanueva, L. E. (2002).** *Fusobacterium nucleatum* in endodontic flare-ups. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* **93**, 179–183.
- Clarke, J. H. (1999).** Toothaches and death. *J Hist Dent* **47**, 11–13.
- Corsten, M. J., Shamji, F. M., Odell, P. F., Frederico, J. A., Laframboise, G. G., Reid, K. R., Vallieres, E. & Matzinger, F. (1997).** Optimal treatment of descending necrotising mediastinitis. *Thorax* **52**, 702–708.
- Coticchia, J. M., Getnick, G. S., Yun, R. D. & Arnold, J. E. (2004).** Age-, site-, and time-specific differences in pediatric deep neck abscesses. *Arch Otolaryngol Head Neck Surg* **130**, 201–207.
- Coulthard, M. & Isaacs, D. (1991).** Retropharyngeal abscess. *Arch Dis Child* **66**, 1227–1230.
- Davis, W. M., Jr & Balcom, J. H., III (1969).** Lincomycin studies of drug absorption and efficacy. An evaluation by double-blind technique in treatment of odontogenic infections. *Oral Surg Oral Med Oral Pathol* **27**, 688–696.
- Dodson, T. B., Perrott, D. H. & Kaban, L. B. (1989).** Pediatric maxillofacial infections: a retrospective study of 113 patients. *J Oral Maxillofac Surg* **47**, 327–330.
- Dymock, D., Weightman, A. J., Scully, C. & Wade, W. G. (1996).** Molecular analysis of microflora associated with dentoalveolar abscesses. *J Clin Microbiol* **34**, 537–542.
- Dzink, J. L., Sheenan, M. T. & Socransky, S. S. (1990).** Proposal of three subspecies of *Fusobacterium nucleatum* Knorr 1922: *Fusobacterium nucleatum* subsp. *nucleatum* subsp. nov., comb. nov.; *Fusobacterium nucleatum* subsp. *polymorphum* subsp. nov., nom. rev., comb. nov.; and *Fusobacterium nucleatum* subsp. *vincentii* subsp. nov., nom. rev., comb. nov. *Int J Syst Bacteriol* **40**, 74–78.
- Facklam, R. (2002).** What happened to the streptococci? Overview of taxonomic and nomenclature changes. *Clin Microbiol Rev* **15**, 613–630.
- Fazakerley, M. W., McGowan, P., Hardy, P. & Martin, M. V. (1993).** A comparative study of cephadrine, amoxicillin and phenoxymethylpenicillin in the treatment of acute dentoalveolar infection. *Br Dent J* **174**, 359–363.
- Flynn, T. R., Shanti, R. M. & Hayes, C. (2006).** Severe odontogenic infections, part 2: prospective outcomes study. *J Oral Maxillofac Surg* **64**, 1104–1113.
- Fouad, A. F., Rivera, E. M. & Walton, R. E. (1996).** Penicillin as a supplement in resolving the localized acute apical abscess. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* **81**, 590–595.
- Gill, Y. & Scully, C. (1990).** Orofacial odontogenic infections: review of microbiology and current treatment. *Oral Surg Oral Med Oral Pathol* **70**, 155–158.
- Gilmore, W. C., Jacobus, N. V., Gorbach, S. L., Doku, H. C. & Tally, F. P. (1988).** A prospective double-blind evaluation of penicillin versus clindamycin in the treatment of odontogenic infections. *J Oral Maxillofac Surg* **46**, 1065–1070.
- Gomes, B. P., Jacinto, R. C., Pinheiro, E. T., Sousa, E. L., Zaia, A. A., Ferraz, C. C. & Souza-Filho, F. J. (2006).** Molecular analysis of *Filifactor alocis*, *Tannerella forsythia*, and *Treponema denticola* associated with primary endodontic infections and failed endodontic treatment. *J Endod* **32**, 937–940.
- Gorbach, S. L., Gilmore, W. C., Jacobus, N. V., Doku, H. C. & Tally, F. P. (1991).** Microbiology and antibiotic resistance in odontogenic infections. *Ann Otol Rhinol Laryngol Suppl* **154**, 40–42.
- Goumas, P. D., Naxakis, S. S., Papavasiliou, D. A., Moschovakis, E. D., Tsintzos, S. J. & Skoutelis, A. (1997).** Periapical abscesses: causal bacteria and antibiotic sensitivity. *J Chemother* **9**, 415–419.
- Graham, D. B., Webb, M. D. & Seale, N. S. (2000).** Pediatric emergency room visits for nontraumatic dental disease. *Pediatr Dent* **22**, 134–140.
- Han, J. K. & Kerschner, J. E. (2001).** *Streptococcus milleri*: an organism for head and neck infections and abscess. *Arch Otolaryngol Head Neck Surg* **127**, 650–654.

- Hanna, C. B., Jr (1991).** Cefadroxil in the management of facial cellulitis of odontogenic origin. *Oral Surg Oral Med Oral Pathol* **71**, 496–498.
- Heimdahl, A., von Konow, L., Satoh, T. & Nord, C. E. (1985).** Clinical appearance of orofacial infections of odontogenic origin in relation to microbiological findings. *J Clin Microbiol* **22**, 299–302.
- Information & Statistics Division (2007).** NHS National Services Scotland. Reference ISD/HIG/IR2005-01474. Edinburgh, UK.
- Jacinto, R. C., Gomes, B. P., Shah, H. N., Ferraz, C. C., Zaia, A. A. & Souza-Filho, F. J. (2006).** Incidence and antimicrobial susceptibility of *Porphyromonas gingivalis* isolated from mixed endodontic infections. *Int Endod J* **39**, 62–70.
- Jimenez, Y., Bagan, J. V., Murillo, J. & Poveda, R. (2004).** Odontogenic infections. Complications. Systemic manifestations. *Med Oral Patol Oral Cir Bucal* **9** (Suppl.), 143–147.
- Khemaleelakul, S., Baumgartner, J. C. & Pruksakorn, S. (2002).** Identification of bacteria in acute endodontic infections and their antimicrobial susceptibility. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* **94**, 746–755.
- Kolokotronis, A. (1999).** Beta-lactamases producing anaerobic bacteria in dentoalveolar abscesses. *J Oral Sci* **41**, 187–190.
- Kulekci, G., Inanc, D., Kocak, H., Kasapoglu, C. & Gumru, O. Z. (1996).** Bacteriology of dentoalveolar abscesses in patients who have received empirical antibiotic therapy. *Clin Infect Dis* **23** (Suppl. 1), S51–S53.
- Kuriyama, T., Karasawa, T., Nakagawa, K., Saiki, Y., Yamamoto, E. & Nakamura, S. (2000a).** Bacteriologic features and antimicrobial susceptibility in isolates from orofacial odontogenic infections. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* **90**, 600–608.
- Kuriyama, T., Nakagawa, K., Karasawa, T., Saiki, Y., Yamamoto, E. & Nakamura, S. (2000b).** Past administration of beta-lactam antibiotics and increase in the emergence of beta-lactamase-producing bacteria in patients with orofacial odontogenic infections. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* **89**, 186–192.
- Kuriyama, T., Karasawa, T., Nakagawa, K., Yamamoto, E. & Nakamura, S. (2001).** Incidence of beta-lactamase production and antimicrobial susceptibility of anaerobic gram-negative rods isolated from pus specimens of orofacial odontogenic infections. *Oral Microbiol Immunol* **16**, 10–15.
- Kuriyama, T., Karasawa, T., Nakagawa, K., Nakamura, S. & Yamamoto, E. (2002a).** Antimicrobial susceptibility of major pathogens of orofacial odontogenic infections to 11 beta-lactam antibiotics. *Oral Microbiol Immunol* **17**, 285–289.
- Kuriyama, T., Karasawa, T., Nakagawa, K., Yamamoto, E. & Nakamura, S. (2002b).** Bacteriology and antimicrobial susceptibility of gram-positive cocci isolated from pus specimens of orofacial odontogenic infections. *Oral Microbiol Immunol* **17**, 132–135.
- Kuriyama, T., Absi, E. G., Williams, D. W. & Lewis, M. A. (2005).** An outcome audit of the treatment of acute dentoalveolar infection: impact of penicillin resistance. *Br Dent J* **198**, 759–763.
- Kuriyama, T., Karasawa, T., Williams, D. W., Nakagawa, K. & Yamamoto, E. (2006).** An increased prevalence of beta-lactamase-positive isolates in Japanese patients with dentoalveolar infection. *J Antimicrob Chemother* **58**, 708–709.
- Lewis, M. A., McGowan, D. A. & MacFarlane, T. W. (1986).** Short-course high-dosage amoxicillin in the treatment of acute dentoalveolar abscess. *Br Dent J* **161**, 299–302.
- Lewis, M. A., MacFarlane, T. W. & McGowan, D. A. (1989).** Antibiotic susceptibilities of bacteria isolated from acute dentoalveolar abscesses. *J Antimicrob Chemother* **23**, 69–77.
- Lewis, M. A., MacFarlane, T. W. & McGowan, D. A. (1990).** A microbiological and clinical review of the acute dentoalveolar abscess. *Br J Oral Maxillofac Surg* **28**, 359–366.
- Lewis, M. A., Carmichael, F., MacFarlane, T. W. & Milligan, S. G. (1993).** A randomised trial of co-amoxiclav (Augmentin) versus penicillin V in the treatment of acute dentoalveolar abscess. *Br Dent J* **175**, 169–174.
- Lewis, M. A. O., Parkhurst, C. L., Douglas, C. W. I., Martin, M. V., Absi, E. G., Bishop, P. A. & Jones, S. A. (1995).** Prevalence of penicillin-resistant bacteria in acute suppurative oral infection. *J Antimicrob Chemother* **35**, 785–791.
- Mangundjaja, S. & Hardjwinata, K. (1990).** Clindamycin versus ampicillin in the treatment of odontogenic infections. *Clin Ther* **12**, 242–249.
- Marty-Ane, C. H., Berthet, J. P., Alric, P., Pegis, J. D., Rouviere, P. & Mary, H. (1999).** Management of descending necrotizing mediastinitis: an aggressive treatment for an aggressive disease. *Ann Thorac Surg* **68**, 212–217.
- Mihos, P., Potaris, K., Gakidis, I., Papadakis, D. & Rallis, G. (2004).** Management of descending necrotizing mediastinitis. *J Oral Maxillofac Surg* **62**, 966–972.
- Nair, P. N. (2004).** Pathogenesis of apical periodontitis and the causes of endodontic failures. *Crit Rev Oral Biol Med* **15**, 348–381.
- Palmer, N. O. A., Martin, M. V., Pealing, R. V. & Ireland, R. S. (2000).** An analysis of antibiotic prescriptions from general dental practice in England. *J Antimicrob Chemother* **46**, 1033–1035.
- Paterson, S. A. & Curzon, M. E. (1993).** The effect of amoxycillin versus penicillin V in the treatment of acutely abscessed primary teeth. *Br Dent J* **174**, 443–449.
- Reader, C. M., Boniface, M. & Bujanda-Wagner, S. (1994).** Refractory endodontic lesion associated with *Staphylococci aureus*. *J Endod* **20**, 607–609.
- Riggio, M. P., Aga, H., Murray, C. A., Jackson, M. S., Lennon, A., Hammersley, N. & Bagg, J. (2006).** Identification of bacteria associated with spreading odontogenic infections by 16S rRNA gene sequencing. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* **103**, 610–617.
- Rocas, I. N. & Siqueira, J. F., Jr (2005).** Detection of novel oral species and phylotypes in symptomatic endodontic infections including abscesses. *FEMS Microbiol Lett* **250**, 279–285.
- Roche, Y. & Yoshimori, R. N. (1997).** In-vitro activity of spiramycin and metronidazole alone or in combination against clinical isolates from odontogenic abscesses. *J Antimicrob Chemother* **40**, 353–357.
- Sakamoto, H., Kato, H., Sato, T. & Sasaki, J. (1998).** Semiquantitative bacteriology of closed odontogenic abscesses. *Bull Tokyo Dent Coll* **39**, 103–107.
- Sakamoto, M., Rocas, I. N., Siqueira, J. F., Jr & Benno, Y. (2006).** Molecular analysis of bacteria in asymptomatic and symptomatic endodontic infections. *Oral Microbiol Immunol* **21**, 112–122.
- Sancho, L. M. M., Minamoto, H., Fernandez, A., Sennes, L. U. & Jatene, F. B. (1999).** Descending necrotizing mediastinitis: a retrospective surgical experience. *Eur J Cardiothorac Surg* **16**, 200–205.
- Sassone, L. M., Fidel, R. A., Faveri, M., Guerra, R., Figueiredo, L., Fidel, S. R. & Feres, M. (2008).** A microbiological profile of symptomatic teeth with primary endodontic infections. *J Endod* **34**, 541–545.
- Schuen, N. J., Panzer, J. D. & Atkinson, W. H. (1974).** A comparison of clindamycin and penicillin V in the treatment of oral infections. *J Oral Surg* **32**, 503–505.
- Schuman, N. J. & Turner, J. E. (1999).** The clinical significance of beta hemolytic streptococci of the milleri group in oral abscesses. *J Clin Pediatr Dent* **23**, 137–142.
- Siqueira, J. F., Jr & Rocas, I. N. (2003a).** Detection of *Filifactor alocis* in endodontic infections associated with different forms of periradicular diseases. *Oral Microbiol Immunol* **18**, 263–265.

- Siqueira, J. F., Jr & Rocas, I. N. (2003b).** Positive and negative bacterial associations involving *Dialister pneumosintes* in primary endodontic infections. *J Endod* **29**, 438–441.
- Siqueira, J. F., Jr & Rocas, I. N. (2003c).** *Pseudoramibacter alactolyticus* in primary endodontic infections. *J Endod* **29**, 735–738.
- Siqueira, J. F., Jr & Rocas, I. N. (2004a).** Nested PCR detection of *Centipeda periodontii* in primary endodontic infections. *J Endod* **30**, 135–137.
- Siqueira, J. F., Jr & Rocas, I. N. (2004b).** Simultaneous detection of *Dialister pneumosintes* and *Filifactor alocis* in endodontic infections by 16S rDNA-directed multiplex PCR. *J Endod* **30**, 851–854.
- Siqueira, J. F., Jr & Rocas, I. N. (2004c).** *Treponema* species associated with abscesses of endodontic origin. *Oral Microbiol Immunol* **19**, 336–339.
- Siqueira, J. F., Jr & Rocas, I. N. (2005).** Uncultivated phylotypes and newly named species associated with primary and persistent endodontic infections. *J Clin Microbiol* **43**, 3314–3319.
- Siqueira, J. F., Jr & Rocas, I. N. (2006).** *Catonella morbi* and *Granulicatella adiacens*: new species in endodontic infections. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* **102**, 259–264.
- Siqueira, J. F., Jr, Rocas, I. N., Oliveira, J. C. & Santos, K. R. (2001a).** Detection of putative oral pathogens in acute periradicular abscesses by 16S rDNA-directed polymerase chain reaction. *J Endod* **27**, 164–167.
- Siqueira, J. F., Jr, Rocas, I. N., Oliveira, J. C. & Santos, K. R. (2001b).** Molecular detection of black-pigmented bacteria in infections of endodontic origin. *J Endod* **27**, 563–566.
- Siqueira, J. F., Jr, Rocas, I. N., Favieri, A., Oliveira, J. C. & Santos, K. R. (2001c).** Polymerase chain reaction detection of *Treponema denticola* in endodontic infections within root canals. *Int Endod J* **34**, 280–284.
- Siqueira, J. F., Jr, Rocas, I. N., Souto, R., Uzeda, M. & Colombo, A. P. (2001d).** Microbiological evaluation of acute periradicular abscesses by DNA-DNA hybridization. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* **92**, 451–457.
- Siqueira, J. F., Rocas, I. N., De Uzeda, M., Colombo, A. P. & Santos, K. R. (2002a).** Comparison of 16S rDNA-based PCR and checkerboard DNA-DNA hybridisation for detection of selected endodontic pathogens. *J Med Microbiol* **51**, 1090–1096.
- Siqueira, J. F., Jr, Rocas, I. N., Souto, R., De Uzeda, M. & Colombo, A. P. (2002b).** *Actinomyces* species, streptococci, and *Enterococcus faecalis* in primary root canal infections. *J Endod* **28**, 168–172.
- Siqueira, J. F., Jr, Rocas, I. N., Andrade, A. F. & De Uzeda, M. (2003).** *Peptostreptococcus micros* in primary endodontic infections as detected by 16S rDNA-based polymerase chain reaction. *J Endod* **29**, 111–113.
- Siqueira, J. F., Jr, Jung, I. Y., Rocas, I. N. & Lee, C. Y. (2005).** Differences in prevalence of selected bacterial species in primary endodontic infections from two distinct geographic locations. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* **99**, 641–647.
- Smith, A. J. & Jackson, M. S. (2003).** Susceptibility of viridans group streptococci isolated from dento-alveolar infections to eight antimicrobial agents. *J Antimicrob Chemother* **52**, 1045–1046.
- Smith, A. J., Jackson, M. S. & Bagg, J. (2001).** The ecology of *Staphylococcus* species in the oral cavity. *J Med Microbiol* **50**, 940–946.
- Storoe, W., Haug, R. H. & Lillich, T. T. (2001).** The changing face of odontogenic infections. *J Oral Maxillofac Surg* **59**, 739–748.
- Sundqvist, G., Johansson, E. & Sjögren, U. (1989).** Prevalence of black-pigmented bacteroides species in root canal infections. *J Endod* **15**, 13–19.
- Tan, P. T., Chang, L. Y., Huang, Y. C., Chiu, C. H., Wang, C. R. & Lin, T. Y. (2001).** Deep neck infections in children. *J Microbiol Immunol Infect* **34**, 287–292.
- Thomas, S. J., Atkinson, C., Hughes, C., Revington, P. & Ness, A. R. (2008).** Is there an epidemic of admissions for surgical treatment of dental abscesses in the UK? *BMJ* **336**, 1219–1220.
- Tung-Yiu, W., Jehn-Shyun, H., Ching-Hung, C. & Hung-An, C. (2000).** Cervical necrotizing fasciitis of odontogenic origin: a report of 11 cases. *J Oral Maxillofac Surg* **58**, 1347–1352.
- Turner Thomas, T. (1908).** Ludwig's angina. An anatomical, clinical, and statistical study. *Ann Surg* **47**, 161–163.
- Valle, G., Quirós, L. M., Andrés, M. T. & Fierro, J. F. (1998).** A  $\beta$ -lactamase belonging to group 2e from oral clinical isolates of *Prevotella intermedia*. *FEMS Microbiol Lett* **158**, 191–194.
- Van Reenan, J. F. & Coogan, M. M. (1970).** Clostridia isolated from human mouths. *Arch Oral Biol* **15**, 845–848.
- von Konow, L., Kondell, P. A., Nord, C. E. & Heimdahl, A. (1992).** Clindamycin versus phenoxymethylpenicillin in the treatment of acute orofacial infections. *Eur J Clin Microbiol Infect Dis* **11**, 1129–1135.
- Wade, W. G., Lewis, M. A., Cheeseman, S. L., Absi, E. G. & Bishop, P. A. (1994).** An unclassified *Eubacterium* taxon in acute dento-alveolar abscess. *J Med Microbiol* **40**, 115–117.
- Wang, L. F., Kuo, W. R., Tsai, S. M. & Huang, K. J. (2003).** Characterizations of life-threatening deep cervical space infections: a review of one hundred ninety-six cases. *Am J Otolaryngol* **24**, 111–117.
- Wang, J., Ahani, A. & Pogrel, M. A. (2005).** A five-year retrospective study of odontogenic maxillofacial infections in a large urban public hospital. *Int J Oral Maxillofac Surg* **34**, 646–649.