PROCEEDINGS OF THE PATHOLOGICAL SOCIETY OF GREAT BRITAIN AND IRELAND

*The 162nd Meeting of the Society was held at Addenbrooke's Hospital,
University of Cambridge on January 3-5, 1991*

SYNOPSIS OF PAPERS OF MICROBIOLOGICAL INTEREST

SYMPOSIUM: MICROBIAL ECOLOGY IN MIXED POPULATIONS

Chairman: Rosamund J. Williams

IN-VITRO MODELS OF THE ORAL MICROFLORA

P. D. Marsh

Pathology Division, PHLS-CAMR, Salisbury, Wilts, SP4 0JG

The microflora of the mouth is diverse and varies between sites within the oral cavity. It forms part of the host defences by preventing colonisation by exogenous micro-organisms. Although this microflora is generally stable, the organisms associated with teeth (dental plaque) can undergo shifts in their relative proportions that can predispose a site to caries and periodontal diseases. Consequently, several in-vitro model systems have been developed to determine the role of the oral microflora in health (colonisation resistance) and the factors responsible for the transition of this microflora from having a commensal to a pathogenic relationship with the host. These systems fall into two main categories: (1) those which attempt to be models of certain unique aspects of the mouth ("artificial mouths"), and which usually contain relevant surfaces to mimic plaque development on teeth; and (2) those in which the modelling of a specific situation is studied under defined and controlled conditions (e.g., chemostats). Both systems often use habitat-simulating media containing glycoproteins (mucins). These systems have been inoculated with both simple combinations (2-4 species) or complex mixtures (either defined or undefined) of oral bacteria. Various types of "artificial mouth" have been used to study colonisation resistance in "plaque" against implantation by *Streptococcus mutans*, and the interaction of dietary sugars and oral sugar clearance rates on pH changes. Chemostat studies have been able to separate the effects of closely-related variables such as sugar availability and low pH, and have shown that it is the latter that favours the growth of potentially cariogenic bacteria. Likewise, serum and elevated pH have been shown to select for anaerobic gram-negative periodontopathogens in mixed cultures. Both systems have their strengths and weaknesses, and are appropriate for different applications; both have contributed to our understanding of the ecology of the mouth in health and disease.

IN-VIVO STUDIES OF MOUTH FLORA

D. Beighton

Hunterian Dental Research Unit, London Hospital Medical College, Turner Street, London, E1 2AD

The dental plaque flora provides a convenient and readily accessible model for studying bacterial growth *in vivo*. The mean generation time (mgt) of bacteria growing in dental plaque during the first 24 h of plaque formation in rats, monkeys and man is 3-4 h; bacterial growth rates in mature plaque are not known. The mgt of plaque bacteria is unaffected by the availability or composition of the host's diet suggesting that the liberation of nutrients from host-derived salivary components determines the rate of bacterial growth. The production of specific glycosidase activities (α -

fucosidase, sialidase and B-N-acetylglucosaminidase) and proteolytic enzyme activities are elevated in the plaque of fasted monkeys. The glycosidase activities could be modulated by N-acetylglucosamine, but not by glucose, added to the drinking water. Such activities are required for the degradation of glycoproteins whose carbohydrate components are common to all animal species. These studies indicate that bacterial growth *in vivo* requires specific glycosidase and non-specific protease activities which are usually provided by the bacterial flora acting in concert, and less commonly by single bacterial species.

A CHEMOSTAT MODEL OF THE HUMAN GUT AND THE ROLE OF GUT MICROFLORA IN THE BREAKDOWN OF FIBRE

G. T. MacFarlane

*MRC Dunn Clinical Nutrition Centre, 100 Tennis Court Road,
Cambridge, CB2 1QL*

Through the breakdown of carbohydrates and proteins to short chain fatty acids, anaerobic bacteria in the human large intestine play an important role in host digestive processes. Ileal effluent, rich in dietary residues and various gastro-intestinal tract secretions, enters the right colon at the caecum, which is a site of intense microbial activity. After a period, the digesta are passed to the left colon for storage and eventual disposal. Due to fermentation in the right colon, substrate availability and, consequently, bacterial activities are likely to be markedly decreased in the distal bowel. Since the healthy large intestine, and the right colon in particular, are inaccessible for most practical purposes, a 3-stage continuous culture system was developed to reproduce, *in vitro*, the relative conditions of substrate availability, bacterial growth rates and pH found in different regions of the gut. Stable bacterial populations characteristic of those occurring in the colon can be maintained for extended periods in the system which facilitates long-term experiments on the metabolism of gut populations. Studies on the degradation of starch and hemicellulosic polysaccharides, by bacteria growing in the gut model were described.

THE ROLE OF THE INTESTINAL BACTERIAL FLORA IN COLONISATION RESISTANCE IN MAN AND ANIMALS

G. C. Mead

*University of Bristol, Department of Veterinary Medicine,
Langford House, Langford, Bristol, BS18 7DU*

Protection of the host against invading pathogens is an important function of the normal, adult, intestinal microflora, and has been demonstrated for man and for animals as diverse as insects, mice and chickens. This phenomenon is being exploited in the poultry industry to control symptomless carriage of food-poisoning salmonellas in intensive production systems, thereby reducing the subse-

quent risk of food-borne illness. Initial development of defined mixtures of chicken caecal bacteria led to protection of orally dosed chicks against an oral salmonella challenge. However, such mixtures were found to be less protective and less stable in their activity than direct cultures of adult gut material, which are now used with appropriate safeguards to avoid transmission of any disease agents. Successful application to commercial poultry flocks was demonstrated in 20 out of 22 field trials. No adverse effects on bird health, growth performance or feed conversion have been reported. The protective effect in the gut is primarily bacteriostatic, but the mechanism of protection is unknown and likely to be multi-factorial. The effect is rapid and apparently independent initially of the immune system; its scope extends to various food-borne pathogens that may be carried asymptotically in the poultry gut, e.g., *Campylobacter jejuni*, *Clostridium perfringens*, pathogenic *Escherichia coli*.

GREASY PIG DISEASE: INTERACTIONS OF THE SKIN MICROFLORA

D. H. Lloyd

Dermatology Unit, The Royal Veterinary College, Hawkshead Lane, North Mymms, Hatfield, Herts, AL9 7TA

Greasy pig disease or exudative epidermitis is a sporadic disease of young pigs which can cause mortality and stunting.

The causative organism, *Staphylococcus hyicus*, is found on normal pig skin and it is unclear why some pigs develop the disease. Poor management and stress factors have been incriminated. When inoculated on to the skin surface of gnotobiotic piglets, *S. hyicus* invariably causes a very severe disease. We have attempted to control this by bacterial interference. The disease can be considerably ameliorated by the prior inoculation on to the skin of antibiotic-producing bacteria. It can be completely prevented by prior inoculation of an avirulent strain of *S. hyicus*. We have investigated the practicability of establishing an antagonistic flora on pig skin under farming conditions. Studies have shown that the normal flora of the pig is created almost immediately at birth. Changes in the flora of the sow's vulva and vagina immediately before parturition appear to promote this. Once the flora is established it is difficult to introduce other bacteria. Thus, it may be necessary to insert potentially antagonistic organisms into the vulval flora of the sow prior to parturition to ensure that they are inoculated on to, and become established in, the piglet skin flora at the earliest opportunity.

SYMPOSIUM: NEW DEVELOPMENTS IN VACCINES

Chairman: E. Mary Cooke

IMMUNITY TO SALMONELLAE AND POTENTIAL VACCINES

Carlos E. Hormaeche

Cambridge University Department of Pathology, Tennis Court Road, Cambridge, CB2 1QP

Live attenuated salmonella vaccines are potent immunogens and are candidate vaccines against typhoid fever and animal salmonellosis. Aro mutants, carrying defined deletions in genes of the aromatic pathway, grow very slowly in the tissues due to their requirement for PABA which is unavailable in mammalian tissues. The virulence of Aro mutants is not markedly increased in moderately immunosuppressed mice (sublethal irradiation or the *xid* defect), suggesting that they may not be a major hazard for immunocompromised populations. Protection probably requires a cellular immune response, which may not be directed against the main LPS O-antigen. Recombinant antigens from viruses, bacteria, toxins and parasites have been expressed in salmonellae as combined vaccines, and protection from disease has been demonstrated with combined vaccines for influenza, streptococci, tetanus and malaria. These constructs show great potential as combined oral vaccines.

PERTUSSIS VACCINE UPDATE

M. J. Corbel

Division of Bacteriology, NIBSC, Blanche Lane, South Mimms, Potters Bar, Herts, EN6 3QG

Current whole-cell vaccines against pertussis have proved effective in controlling the disease but concern over their reactogenicity has stimulated the search for acellular

vaccines of more precisely defined composition. Two main approaches have been adopted: the formulation of vaccines enriched in respect of putative major protective antigens and depleted of reactogenic components, especially endotoxin and active pertussis toxin (PT) (Takeda type vaccines); and vaccines prepared from purified individual protective antigens (Biken type). Emphasis has been placed on the inclusion of detoxified PT and filamentous haemagglutinin (FHA) as the main components of acellular vaccines. However, these alone may not ensure adequate immunity and additional components, including agglutinogens 2 and 3 and the 69-Kda outer-membrane protein (pertactin), have been added to some formulations. The detoxification of PT still presents technical problems and alternatives to toxoiding by cross-linking reagents have been explored. These include the use of synthetic peptides corresponding to major epitopes of PT sub-units and the production of genetically detoxified toxin protein. There is a lack of data on the optimum composition of acellular pertussis vaccines which can only be obtained from clinical trials which include the whole-cell vaccine.

THE PROSPECTS FOR AN AIDS VACCINE

P. J. Greenaway

Division of Pathology, Centre for Applied Microbiology and Research, Porton Down, Salisbury, Wilts, SP4 0JG

The control of AIDS is dependent on the development of effective vaccines for use in either selective or mass immunisation programmes. These vaccines should be safe, stable and capable of inducing a long-lasting immunity that will protect individuals from infection by the human immunodeficiency virus. The vaccine finally adopted for

this purpose is likely to be produced by genetic manipulation techniques and based on subunit proteins encoded by HIV. Because of the restricted host range of HIV and the difficulties of mounting adequate clinical trials much of the developmental work will depend on the use of suitable animal model challenge systems. The infection of macaques with strains of the simian immunodeficiency virus, which produces a disease with a pathology similar to that of human AIDS, represents the best animal model currently available. This model has been used to assess crude inactivated vaccines based on either infected cells or purified virus, and protection from both homologous and heterologous live virus challenge has been demonstrated. Studies aimed at investigating the parameters of protection and at refining these crude vaccines are in progress.

IMMUNITY IN HUMAN SCHISTOSOMIASIS: PROSPECTS FOR VACCINATION?

A. E. Butterworth

Department of Pathology, University of Cambridge, Tennis Court Road, Cambridge, CB2 1QP

Safe and effective drugs for the treatment of the three main species of schistosome that affect man have now been available for almost 20 years, and most operational control programmes rely on some form of mass or targeted chemotherapy at the community level. Such programmes, properly delivered, can be useful as a short-term measure. However, they have serious drawbacks, in particular the costs of both drugs and delivery of treatment, and the need for repeated surveillance and re-treatment. Alternative approaches, especially vaccination, would therefore be useful. Studies in experimental animal models have demonstrated the existence of acquired immunity, and have led to the molecular cloning of various candidate vaccine antigens. However, results from these studies cannot be extrapolated directly to man. Summarising the results of a 10-year study of human schistosomiasis *mansoni* in Kenya, we have demonstrated the very slow development, with age, of acquired immunity to re-infection after treatment. We have also obtained evidence that the continued susceptibility of young children is attributable to high levels of "blocking" antibodies, especially IgM, whereas IgE antibodies may contribute to protection in older individuals. The finding

that man does develop immunity to infection bodes well for the eventual development of a vaccine; however, its success will depend on our ability to elicit appropriate immune responses.

VARICELLA VACCINE FOR THE IMMUNOCOMPROMISED

H. Bogaerts

Smith Kline Beecham Biologicals, 89, Rue de l'Institut B-1330, Rixensart, Belgium

Varicella infection is potentially life threatening for children who are immunodeficient or on immuno-suppressive therapy. Visceral dissemination of the virus is frequent and the mortality rate for immunodepressed patients is about 100-fold that of otherwise healthy children. For these reasons, a live attenuated varicella vaccine, containing the Oka viral strain, was developed in Japan in the early 1970s. Studies have shown that this vaccine is safe and immunogenic and that, even if there is some excretion of the vaccine viral strain, there is no reversion to virulence. Although the vaccine does not prevent all varicella infections in the immunocompromised patient, it does protect against the serious complications of the disease. In addition, vaccinated children with leukaemia have not shown a higher incidence of zoster with respect to matched healthy controls. The Oka strain was licensed to a European manufacturer in 1978. Results with this vaccine in approximately 1000 subjects have confirmed previous findings on its safety and immunogenicity. It is the only vaccine registered outside Japan for administration to high risk patients (after interruption of maintenance chemotherapy) and their contacts and for the termination or prevention of epidemics in closed communities. Chicken-pox is often considered a benign disease. However, certain infrequent complications such as encephalitis, Reye's syndrome and bacterial superinfection have been observed, which have raised the issue of the vaccination of healthy children. Cost-benefit studies have demonstrated the advantages of administering the vaccine to all children. This will be most feasible by combining varicella vaccine with measles-mumps-rubella vaccine, although potential problems, i.e., vaccine stability and competition between viral strains, need to be addressed.

ORAL PRESENTATIONS

EFFECT OF NUTRITION ON VACCINATION AGAINST DERMATOPHILOSIS IN AFRICAN CATTLE

Anna B. Sanders and D. H. Lloyd

Dermatology Unit, Royal Veterinary College, Hawkshead Lane, North Mimms, Herts, AL9 7TA

Dermatophilosis is a severe and sometimes fatal bacterial skin disease of ruminants. Although ubiquitous, its worst effects are seen in the humid tropics. In temperate climates the disease does not cause such a degree of morbidity or mortality. Poor nutrition has been shown to delay the healing of lesions and reduce the efficacy of vaccines against *Dermatophilus congolensis*, the causative organism. This study aimed to investigate the effect of supplementary feeding on the effectiveness of vaccination in Tchad, where

malnutrition is endemic in livestock during the dry season. Forty cattle were divided into wellnourished (hay with cottonseed) and malnourished (hay only) groups. Half of each group was given three inoculations of a live bacterial vaccine in Freund's incomplete adjuvant, over a 6-month period. Two weeks after the final vaccination, the animals were challenged with *Dermatophilus* by skin inoculation. The results showed that, in cattle whose diet was supplemented, the vaccines reduced the severity and duration of the experimental lesions; whereas in animals without the dietary supplements these effects were much less marked. The study also demonstrated that animals whose skin was unpigmented had more severe lesions. This effect may have been due to the interaction of solar radiation and unpigmented skin.

MOLECULAR CLONING OF AN *ONCHOCERCA VOLVULUS* ANTIGEN THAT SHOWS IMMUNOLOGICAL CROSS-REACTIVITY WITH A 40 000-KDA COMPONENT OF RETINAL PIGMENT EPITHELIUM

Gabriele Braun, V. Connor, M. D. Downham, F. Engelbrecht, C. Gilbert, N. M. McKechnie, J. A. Whitworth and D. W. Taylor

Department of Pathology, Tennis Court Road, Cambridge, CB2 1QP

Onchocerca volvulus is the causative agent of river blindness or onchocerciasis. Infection with *O. volvulus* is often associated with severe skin disease, visual impairment and blindness. It has been suggested that eye pathology, including chorioretinopathy, may result from autoimmune responses. We have used Western blotting techniques and immunocytochemistry, in addition to ELISA, to investigate possible cross-reactivity between *O. volvulus* and eye tissue. An antiserum raised against a 22 000-Kda recombinant antigen isolated from an expressing adult *O. volvulus* cDNA library, recognised a 44 000-Kda antigen of bovine retinal pigment epithelium (RPE). The same antiserum used in immunocytochemical studies localised the target antigen to cytoplasm of human RPE. The *O. volvulus* cDNA clone has been sequenced but no homology has been found with any human tissue.

MOLECULAR, IMMUNOLOGICAL AND BIOLOGICAL CHARACTERISATION OF A TOXIN A-NEGATIVE TOXIN B-POSITIVE STRAIN OF *CLOSTRIDIUM DIFFICILE*

S. P. Borriello, S. Hyde, S. V. Seddon, B. W. Wren,* M. M. Krishna, S. Tabaqchali* and S. Manek†

*Microbial Pathogenicity Research Group, Clinical Research Centre, Watford Road, Harrow, Middlesex, *Department of Medical Microbiology, St Bartholomew's Hospital, London and †Department of Histopathology, Northwick Park Hospital, Harrow*

A cytotoxic strain of *Clostridium difficile* was shown to be negative for toxin A by ELISA with the monoclonal antibody PCG-4 (Strain A-B+). Molecular analysis demonstrated that the lack of reaction with PCG-4 was not due to a problem of translation but absence of that part (3' end) of the toxin A gene. DNA probes for the 5' end of the gene indicated that much, if not all, of this part of the gene was also absent. The toxin B of this strain eluted at exactly the same point as toxin B from a conventional fully toxigenic strain (VPI 10463) by fast protein liquid chromatography, but at a point different from that of the toxin B equivalent (LT) of *C. sordellii*. Antibody raised to the toxin B of strain A-B+ neutralised the cytotoxic activity of toxin B from strain VPI 10463 but not that of toxin A. Conversely, antibodies to strain VPI 10463, as well as *C. sordellii*, neutralised the cytotoxicity of the A-B+ strain. This strain also produced proteolytic and hydrolytic enzymes. Administration of this strain to hamsters 5 d after dosing with clindamycin resulted in wet-tail and death starting about 4 d after challenge with *C. difficile*, unlike controls which received a fully toxigenic strain and died the next day. However, delayed death appears to be a feature of increased sensitivity to clindamycin, as following challenge with the A-B+ strain 10 d post clindamycin resulted in death within 48 h. Histological analysis showed that there was some small bowel damage, tufting of the surface epithelium, and haemorrhage in the caecum, with mild crypt hyperplasia and increased mitotic activity.

PURIFICATION AND CHARACTERISATION OF A CELL-ASSOCIATED PROTEASE FROM *CLOSTRIDIUM DIFFICILE*

S. V. Seddon and S. P. Borriello

Microbial Pathogenicity Research Group, Clinical Research Centre, Watford Road, Harrow, Middlesex HA1 3UJ

A cell-associated protease was purified from a known highly virulent toxigenic strain of *Clostridium difficile*. Cells were sonicated, centrifuged and the resulting supernate was passed through Sepharose QFF and Mono Q FPLC anion exchange columns. Positive fractions were determined in a microtitration plate assay with the synthetic substrate N α -Benzoyl-D.L arginine to nitroanilide with an ELISA plate reader at 412 nm. Final purification was achieved with a Superose 12 size exclusion FPLC column. The purification procedure resulted in a 14-fold increase in purity with a 10% yield. The mol. wt was determined to be c. 45 Kda on native PAGE and the pI value to be 4.5 with the Pharmacia Phast system. The purified enzyme had temperature and pH optima of 37°C and 7.5 respectively. Substrate specificity and inhibition susceptibility suggested that the enzyme was similar to the protease clostripain expressed by *C. histolyticum*. The possible role of this enzyme in the metabolism and pathogenesis of *C. difficile* is unclear and will require further study.

FLOW CYTOMETRY AS A METHOD OF ASSESSING BACTERIAL PHAGOCYTOSIS AND INTRACELLULAR KILLING

Carol A. Webster and R. C. B. Slack

Department of Microbiology and PHLS, University Hospital, Queens' Medical Centre, Nottingham, NG7 2UH

Flow cytometry as a technique has been in use for 40 years. The introduction of laser beam illumination and computerised graphics has meant that an extremely sophisticated technique has become routinely available. We have used flow cytometry to assess phagocytosis and intracellular killing for which conventional methods are labour intensive and generally unreliable. In addition it is often only one aspect of the process which is investigated. However, by using a 'FacsCan' flow cytometer, employing a combination of fluorescent dyes and cellular light scatter, the whole process can be monitored quickly and accurately, the sensitivity of the machine being 10^2 organisms/ml. We used both human polymorphonuclear neutrophils and mouse resident peritoneal macrophages with a clinical strain of *Staphylococcus aureus* stained with fluorescein isothiocyanate. Phagocytosis was monitored with phagocyte fluorescence, increasing with the number of attached and internalised bacteria. Attached bacteria can be differentiated by quenching with Trypan blue. Intracellular killing was then measured by releasing the bacteria and staining with propidium iodide, a dye which binds to DNA but which is not taken up by viable cells. Conventional plate counts were performed concurrently, resulting in good correlation. The method is ideal for assessing the effects of immunopotentiating drugs.

EVALUATION OF THE E-TEST, A NOVEL METHOD FOR DETERMINATION OF MICs OF ANTIBIOTICS

D. F. J. Brown and Lynda Brown

Clinical Microbiology and Public Health Laboratory, Addenbrooke's Hospital, Hills Road, Cambridge, CB2 2QW

The E-Test (AB Biodisk) is a diffusion method for measuring minimum inhibitory concentrations (MICs) of bacteria and consists of 50 mm $\times$ 5 mm dry strips carrying

an antibiotic gradient. We have evaluated the performance of the technique by comparison with a conventional agar dilution MIC method. A total of 1304 tests was done on a wide variety of organisms, which were tested against benzyl penicillin, ampicillin, piperacillin, ceftazidime, cefotaxime, gentamicin, amikacin, erythromycin, trimethoprim and ciprofloxacin. Agreement between the two methods was good in that 98.9% of MICs were within two two-fold dilution steps and 86% were within one two-fold dilution step of each other. Of the 15 test results with MICs differing by more than two two-fold dilution steps, 13 were obtained with benzyl penicillin and penicillinase-producing staphylococci. With these organisms, the E-test gave a more reliable indication of resistance mediated by penicillinase than the agar dilution MIC method. The E-test proved to be a reliable method for determining MICs. The test is very rapid to set up and no special equipment is needed. The strips are stored with a dessicant and have a shelf life which is at least as long as paper disks containing antibiotics. The speed and ease with which tests can be set up and read make the system an attractive alternative to conventional MIC methods.

SPECIFIC BINDING OF ATP TO *CLOSTRIDIUM DIFFICILE* TOXIN A

M. D. Lobban, M. Krishna* and S. P. Borriello*

Department of Biochemistry, University of Glasgow, University Avenue, Glasgow, G12 8QQ and *Microbial Pathogenicity Research Group, Clinical Research Centre, Watford Road, Harrow, Middlesex, HA1 3UJ

Clostridium difficile produces two major protein toxins, toxins A and B, both of which are involved in the pathogenesis of gastrointestinal disease caused by this organism. The molecular mechanism of action of these toxins remains unclear. In order to attempt to clarify the mechanisms of action and structure-function relationships, we are trying to identify functions (as distinct from effects). One such function, binding of polyphosphates by toxin B, has been shown by others. Because of the high level of homology between the nucleotide sequences for the toxin A and B genes, we considered it useful to screen toxin A for polyphosphate binding. To achieve this we monitored the quenching of the intrinsic fluorescence of toxin A by ATP at 340 nm following excitation at 280 nm. Titrations were performed with 5- μ l increments of 1 mM ATP yielding a range of 1–200 μ M ATP. A double reciprocal plot showed binding of ATP to be specific and saturated at 30–40 μ M ATP. The results demonstrated that toxin B, like toxin A, also had a phosphate-binding site. This is likely to be coded for by homologous segments of the respective genes and is likely to reside at the 5' end of the molecule.

A HOSPITAL OUTBREAK OF INFECTION WITH *BACILLUS* SPP. ASSOCIATED WITH A BUILDING SITE

Stephanie J. Dancer

The Department of Medical Microbiology, St Bartholomew's Hospital, West Smithfield, London, EC1A 7BE

Over a period of 4 months, an outbreak of infection with *Bacillus* spp. occurred in a London hospital involving patients in the Intensive Care Unit (I.C.U.) and infants on post-natal wards. Sporadic cases were also identified from other parts of the hospital. Forty patients were included in the outbreak, producing isolates from skin and wound sites, arterial lines and blood. The outbreak began following initiation of building works in close proximity to the affected wards. *Bacillus* strains were identified from the construction

site, air-conditioning inlet filters and from all air-vents and settle plates in I.C.U. Transmission of spores between building site, wards and filters was investigated with theatrical smoke bombs. A further source of *Bacillus* spp. was provided by the hospital laundry, also located near the building works. The organism was found from the rollers of two steam-pressers, the floor below, and lint-extract compartments beneath several tumble-driers. Impression plates of laundry directly from steam-pressers and linen stored on the wards also produced *Bacillus* spp. API system tests identified most isolates as *B. cereus* and serotyping established two patient clusters. The combination of summer weather, building works and geographical positioning potentiated this outbreak; whether by laundry contamination or air-borne dissemination, bacillus spores settled directly on to those patients least likely to cope with infection.

IDENTIFICATION OF COMMON PROTEINS IN *HAEMOPHILUS DUCREYI*

Shaula Reilly and Catherine A. Ison

Department of Medical Microbiology, St Mary's Hospital Medical School, Paddington, London, W2 1PG

The protein profiles of whole-cell lysates and outer-membrane preparations of clinical isolates of *Haemophilus ducreyi* were examined by SDS-PAGE and probed with pre-immune and immune rabbit serum. The whole-cell lysates of the clinical isolates showed very similar patterns on SDS-PAGE with at least 10 proteins of the same mol. wt common to all strains. In contrast, the outer-membranes showed differing profiles between strains. On Western blotting, the pre-immune serum reacted with epitopes on five proteins in both whole-cell lysates and outer membranes of *H. ducreyi*. The immune serum gave patterns with all of the whole-cell lysates that were indistinguishable. The outer membranes of each of the strains tested reacted in a very similar manner. This contrasts with the SDS-PAGE profiles. Five proteins that are common to *H. ducreyi* and share epitopes that react with this immune serum have been identified; doublets of 86/82 Kda and 62/60 Kda and a single protein of 40 Kda.

NEW NAMES FOR STREPTOCOCCI

G. Colman

Division of Hospital Infection, Central Public Health Laboratory, Colindale Avenue, London, NW9 5HT

The classification of streptococci is in a state of flux. Of Sherman's four divisions, namely pyogenic, viridans, faecal and lactic, the last two now form new genera: *Enterococcus* and *Lactococcus*. The familiar method of classification from below in which properties are sought to define species (pathogenic, saprophytic or epiphytic) from particular hosts or sites is progressively being replaced by classification from above, based on studies of nucleic acids. The use of the latter procedures combined with the strict application of the rules of nomenclature have led to radical proposals for dividing the streptococci. *Streptococcus pyogenes* (group A) and *S. agalactiae* (group B) remain unchanged but it has been suggested that the name *S. dysgalactiae* be applied to strains of group C (*S. equisimilis* and *S. dysgalactiae*), the human strains of group G and those of group L. The human viridans streptococci are becoming more ordered. *S. sanguis* has been divided into three species. *S. mitior* has been replaced by two: *S. oralis* and *S. mitis*. *S. milleri* can now probably be subdivided usefully into *S. anginosus*, *S. intermedius* and *S. constellatus*. It has, however, been suggested that *S. salivarius* forms one species with *S. thermophilus*. Thirteen species

have been proposed among the enterococci and six among the lactococci. Attempts to produce a phylogenetic classification will probably introduce further changes.

CIPROFLOXACIN'S SLOW RATE OF KILL AGAINST *ENTEROCOCCUS FAECALIS*

I. Morrissey, C. S. Lewin* and J. T. Smith

*Microbiology Section, Department of Pharmaceutics, School of Pharmacy, 29 Brunswick Square, London, WC1N and *Department of Medical Microbiology, University of Edinburgh Medical School, Teviot Place, Edinburgh, EH8 9AG*

MIC determinations indicate that ciprofloxacin is less active against *Enterococcus faecalis* than it is against staphylococci or enterobacteria. However, such determinations do not provide any information on the lethality of the 4-quinolones. Therefore the bactericidal activity of ciprofloxacin at concentrations from 0.15 to 300 mg/L was investigated against *E. faecalis*, *Staphylococcus aureus* and *S. epidermidis* over 24 h in nutrient broth at 37°C. Ciprofloxacin was rapidly bactericidal against *S. aureus* E3T and *S. epidermidis* SK360, killing >99% of the bacteria during the first 3 h of exposure at its optimum bactericidal concentration. A reduction in viability of log₁₀ or less was observed over exposure for the next 21 h. In contrast, ciprofloxacin displayed a much slower rate of kill against *E. faecalis* as little or no bactericidal activity was detected in nutrient broth over the first 3 h for *E. faecalis* ACTC 1943 and two clinical isolates at concentrations up to 150× the MIC. Ciprofloxacin was bactericidal against the enterococci after exposure for 6 h, killing up to 95% of the bacteria. At 24 h, the amount of kill obtained with the 4-quinolone against the enterococci was similar to that obtained against the staphylococci. In conclusion, ciprofloxacin exhibits a reduced rate of kill against *E. faecalis* as well as being less active in terms of MIC. However, as the amount of kill obtained after exposure to the 4-quinolone for 24 h was similar for *E. faecalis* or staphylococci, ciprofloxacin will still be bactericidal against the former if the period of exposure to drug concentrations above the MIC is long enough. These requirements for lethality are met in urine where ciprofloxacin may be used to treat enterococcal UTIs.

ANTIBIOTIC RESISTANCE PLASMIDS IN *STAPHYLOCOCCUS HYICUS*

Louise Kent, Mary Rahman and W. C. Noble

Department of Microbial Diseases, Institute of Dermatology, St Thomas' Hospital, Lambeth Palace Road, London, SE1 7EH

Staphylococcus hyicus strains isolated from normal pigs and pigs with exudative epidermitis are frequently resistant to antibiotics. We report here studies on plasmids which mediate resistance to tetracycline, streptomycin and streptomycin plus chloramphenicol. Some plasmids that mediate resistance to tetracycline clearly belonged to the family of such plasmids reported from staphylococci of human origin, others did not. Streptomycin resistance plasmids all showed the relaxation phenomenon described for strains of human origin and had similar restriction patterns. The streptomycin plus chloramphenicol plasmid also belongs to the same group. This work extends our knowledge of the pool of plasmids which staphylococci share.

INFLUENCE OF SALT CONCENTRATION ON THE DETECTION OF METHICILLIN RESISTANCE IN COAGULASE-NEGATIVE STAPHYLOCOCCI

D. Law, G. M. Megson, Maeve G. L. Keaney and Leela A. Ganguli

Microbiology Department, Hope Hospital, Salford, M6 8HD

Three hundred strains of coagulase-negative staphylococci were screened for methicillin resistance (MR) by the breakpoint method with Columbia agar containing methicillin 4 mg/L and varying concentrations of salt; all were incubated at 30°C. In 68 strains, MR was dependent on salt concentration; some of these strains were MR at low salt concentrations but sensitive at high concentrations whereas others showed the opposite effect. No one single salt concentration was able to correctly identify all of these strains as MR. To detect all MR strains, Columbia agar with added salt 4% and without salt is needed. The use of other media or oxacillin proved inferior. By disk sensitivity test, some MR strains gave very large zones to methicillin and would be classified as sensitive. Varying the salt concentration of the medium had a marked effect on zone size depending on the species of *Staphylococcus* tested. In some cases, target zones were seen at high salt concentrations. Oxacillin gave smaller zones than methicillin but some of the MR strains showed oxacillin zones large enough to be classed as sensitive. For disk testing, the use of oxacillin disks on media without salt and with salt 5% is optimum for detecting MR, a zone of <15 mm on either media indicates methicillin resistance.

CYTOCHEMICAL PROBES OF BACTERIAL VIABILITY

Susan M. Thom and M. R. Barer

Department of Microbiology, The Medical School, Framlington Place, Newcastle upon Tyne, NE2 4HH

In view of the time delay and bias introduced by culture methods, direct determination of the numbers, identities and antimicrobial susceptibilities of the bacteria present in clinical samples is desirable. Cytochemical techniques may be suitable for this purpose. The ability of six cytochemical probes to discriminate between viable and non-viable bacteria was examined as a potential approach to rapid susceptibility testing. Bacteria were immobilised on glass microscope slides and stained with, acridine orange, fluorescein diacetate, ethidium bromide, or a tetrazolium dye (nitroblue tetrazolium, iodonitrotetrazolium or methylthiazoldiphenyl tetrazolium). Light microscopy of stained preparations revealed that only the tetrazolium dyes reliably differentiated viable bacteria from heat- or formaldehyde-treated bacteria. *Staphylococcus aureus* was more intensely stained than *Escherichia coli*. Inclusion of an intermediate electron carrier, such as phenazine methosulphate, accelerated staining such that results could be read after incubation for 20 min at 37°C. Moreover, preliminary results indicate that exponential and stationary-phase cultures may also be differentiated. These studies demonstrate the feasibility of a cytochemical approach to aspects of diagnostic microbiology and suggest that probes of oxidative metabolism, such as the tetrazolium dyes, could be used for the rapid determination of viability.

ULTRASTRUCTURE OF COCCAL FORMS OF *CAMPYLOBACTER JEJUNI*—ARE THESE VIABLE BUT NON-CULTURABLE?

D. M. Jones, A. Curry and E. M. Sutcliffe

Public Health Laboratory, Withington Hospital, Manchester, M20 8LR

On prolonged contact with water, both in the environment

and in laboratory microcosms, the normal curved sinusoid form of *Campylobacter jejuni* is replaced by coccal morphology. In laboratory microcosms, the organisms became non-culturable coccal forms after exposure for 3 weeks to water at 4°C, but many cocci retained the appearance of viability for some months. On negative staining, these cocci appeared to have intact cell membranes and were bi-flagellate with the flagellar filaments arising adjacent to each other and not on opposite poles. By growing *C. jejuni* under stressed conditions the formation of cocci is accelerated and this facilitates the observation of cells intermediate between vibrioid and coccal forms. In negatively-stained preparations, short vibrioid cells appear to bulge laterally from the original convexity and this continues until a sphere is formed. In thin sections, the terminal sub-inner membrane thickenings normally seen at the poles of the sinusoid form are lost. Cocci contain a disk-shaped structure associated with nucleoid filaments, suggesting a dormant state. The cytoplasm and inner membrane do not retain a folded rod morphology seen in coccal forms of *Helicobacter pylori* and this suggests that the cocci of *C. jejuni* are developed in a way quite different from those of that organism.

EFFECT OF IRON LIMITATION ON MENINGOCOCCAL TRANSFER RNA

D. A. Ala'Aldeen, E. Griffiths* and S. P. Borriello

Microbial Pathogenicity Research Group, MRC Clinical Research Centre, Watford Road, Harrow HA1 3UJ and *National Institute for Biological Standards and Control, Balanch Lane, South Mimms, Potters Bar

Neisseria meningitidis expresses several new outer-membrane proteins when grown under iron limitation. The regulation of biosynthesis of these proteins is not understood. In enteric organisms the expression of many iron-regulated genes is negatively controlled by a global repressor protein (fur) which uses ferrous iron as a co-repressor. An additional global regulatory system based on undermodification of several transfer RNAs (tRNA) also appears to operate. Using ³H or ¹⁴C labelled aromatic amino-acids, or both, to generate hot aminoacyl-tRNAs, significant differences were observed between chromatographic elution profiles of tRNA from meningococci grown under iron-sufficient and iron-deficient conditions. In the latter (where desferrioxamine was used as an iron chelator), the majority of tRNA eluted at an ethanol concentration level much lower than the former. Furthermore, addition of EDDA as an iron chelator (in the presence of which meningococci do not grow further but survive and express new iron-regulated proteins) no new tRNA species were detected. These results show that it is growth of the bacteria under stress, rather than mere survival, that encourages the production of the new species of tRNA. This indicates that, when subjected to iron-limitation, the new species of tRNAs are not modified old ones but newly transcribed distinctive species.

A DEMYELINATING VARIANT OF MOUSE HEPATITIS VIRUS CONTAINS A DELETION IN THE ENVELOPE GLYCOPROTEIN AND IS NEURO-ATTENUATED BY ITS REDUCED RATE OF SPREAD

J. K. Fazakerley, Sueanne E. Parker,* F. Bloom* and M. J. Buchmeier*

Department of Pathology, University of Cambridge and *Department of Neuropathology, Scripps Clinic and Research Institute, La Jolla, CA, USA

Intracerebral inoculation of adult mice with mouse hepatitis virus type-4 (MHV4) results in an acute lethal encephalitis. Inoculation with the V5A13.1 monoclonal antibody neutralisation resistant variant of this virus results

in a non-lethal, subacute demyelinating encephalitis. Both viruses replicate in the brain, predominantly in neurones and spread along known neuronal pathways. Certain areas are consistently infected, including the supramammillary, vestibular, arcuate, and deep cerebellar nuclei, the subiculum, hippocampus, anterior olfactory area and the mitral cells of the olfactory lobe. Other areas, including the occipital cortex, the thalamic nuclei and the outer layers of the cerebellum, are rarely infected. Following intranasal infection, both viruses track up the olfactory nerve to infect, sequentially, neurones in the olfactory lobe, the anterior olfactory area and nuclei in the rhinencephalon. The major difference between MHV-4 and its variant is their rate of spread. Parental MHV-4 spreads rapidly, infecting and destroying large numbers of neurones and resulting in death within 4 days of infection. The variant virus spreads more slowly, allowing intervention by the immune response before viral destruction reaches lethal levels. This slow rate of spread is also observed in athymic *nu/nu* mice but, in the absence of an intact immune response, the infection eventually reaches the same extent as MHV-4 resulting in death within 8 days of infection. Direct RNA sequencing of the envelope glycoprotein genes of these two viruses indicates that the variant differs from the parent by a deletion of 142 amino acids in the S1 envelope glycoprotein. It is likely that this hypervariable region determines the rate of viral spread within the CNS and is thus a determinant of neurovirulence.

A SENSITIVE METHOD FOR ISOLATING FUSOBACTERIUM NECROPHORUM FROM FAECES

G. R. Smith and Sarah A. Barton

Institute of Zoology, Regent's Park, London, NW1 4RY

Necrobacillosis occurs in a wide range of domesticated and wild animal species. Lesions of the body surface and mouth are thought to result from faecal contamination of small wounds with virulent *Fusobacterium necrophorum*. However, the isolation of such organisms from heavily contaminated material (e.g., faeces, soil) has been hampered by the lack of an efficient selective medium and by high minimum infective dose. A sensitive method was found to be the subcutaneous injection of faeces, suspended (5% w/v) in broth culture of *Actinomyces (Corynebacterium) pyogenes* or *Staphylococcus aureus* to increase fusobacterial infectivity, into mice pre-treated with clostridial antitoxins. When necrobacillosis developed, the beaded filaments of *F. necrophorum* were identified microscopically in tissue from the advancing edge of the lesion and the organism was isolated on a partly selective medium. The enhancement of fusobacterial infectivity produced by *A. pyogenes* and by *S. aureus* was high, but the latter was slightly more efficient, enabling as few as 80 *F. necrophorum* organisms/g of faeces to be detected. Use of the method showed that 3 of 16 wallabies had *F. necrophorum* in their faeces at the time of examination. Numerous epidemiological applications are suggested.

TYPING OF ORAL CANDIDA ALBICANS ISOLATES FROM AIDS PATIENTS BY MORPHOTYPING, RESISTOTYPING AND RFLP COMPARISON

S. A. Howell, M. A. Pearce, C. Fraser* and J. Midgley†

Department of Microbial Diseases, Institute of Dermatology, St Thomas' Hospital, London, SE1 7EH, *Division of Hospital Infection, Central Public Health Laboratory, Colindale, London, NW9 5HT and †Department of Medical Microbiology, Charing Cross and Westminster Medical School, London, SW1P 2AR

Isolates of *Candida albicans* from AIDS patients with oesophageal candidiasis were obtained over an 18-month

period. Each isolate from every patient was characterised by three methods: morphotyping, resistotyping and comparison of restriction fragment length polymorphisms (RFLP). The first isolation from each patient was obtained prior to antifungal treatment. Subsequent isolates would, therefore, represent recolonisation of the patient with the original strain, or infection with a different strain. The results of RFLP comparisons from a total of 49 isolates from 10 selected patients showed that three patients had carried more than one strain. Ten distinct RFLP types were observed and one type was produced from isolates harboured by 5 of 10 patients. RFLP comparisons were found to be more reproducible than resistotyping and provided greater discrimination than morphotyping.

DIFFERENTIATION BETWEEN SPECIFIC AND NON-SPECIFIC ANTIBODIES TO HEPATITIS C VIRUS IN PATIENTS WITH CHRONIC LIVER DISEASE

J. J. Gray, T. G. Wreghitt, D. G. D. Wight*, V. Sundaresan†, P. J. Friend and R. Y. Calne‡

*Clinical Microbiology and Public Health Laboratory, *Department of Histopathology, and †Department of Surgery, Addenbrooke's Hospital, Hills Road, Cambridge, CB2 2QQ*

The ORTHO Diagnostics System enzyme immunoassay for detecting antibodies to hepatitis C virus (HCV) was used to test serum samples collected from 100 patients with chronic liver disease awaiting liver transplantation. Serum samples from 8 (42%) of 19 patients with cryptogenic cirrhosis were highly reactive (optical density > 1.1; mean 2.38) in the enzyme immunoassay. Samples from a further six patients, with liver disease not normally associated with virus infection, were weakly reactive (OD < 0.9; mean 0.69). To determine whether the low OD values seen with some of the samples were non-specific, all antibody-positive samples were retested with an 8 M urea wash. This was included during the first ELISA wash procedure to dissociate weak-binding, low-avidity antibodies from the antigen. A total of 8 (89%) of 9 HCV antibody-positive patients with cryptogenic cirrhosis had predominantly high-avidity antibodies (< 50% reduction in OD value in the presence of 8 M urea), whereas only 1 (17%) of 6 HCV antibody-positive patients with other chronic liver disease had high-avidity antibody. This may indicate that, in chronic liver disease, there is a group of patients with non-specific low-avidity antibodies that cross-react with the recombinant HCV antigen used in this assay. The interpretation of OD values obtained in this assay, with and without an 8 M urea wash, in conjunction with the results of histological examination of liver biopsy material, may be helpful in distinguishing specific from non-specific reactions.

NALIDIXIC ACID-RESISTANT SALMONELLAS FROM ANIMALS

C. Wray, L. J. V. Piddock* and I. M. McLaren

*Central Veterinary Laboratory, New Haw, Weybridge, Surrey, KT15 3NB and *Antimicrobial Agents Research Group, Department of Medical Microbiology, University of Birmingham, B15 2TJ*

During 1989, 17 nalidixic acid-resistant (nal^R) *Salmonella* serotypes were received from poultry, some of which were imported into the UK and in quarantine. These strains represented six different serotypes and were 0.003% of the total isolates of *Salmonella* from poultry in the UK that year. All the nal^R isolates had a 250-fold increase in the MIC to nalidixic acid compared to the NCTC type strain for each serotype. The nal^R isolates were also less susceptible to newer quinolones that are used clinically in man, such as

ciprofloxacin; for two isolates, the MIC of this agent was equivalent to the UK recommended breakpoint concentration (1 µg/ml). For six of the isolates an MIC ≥ 1 µg/ml (the recommended breakpoint concentration for the veterinary quinolone enrofloxacin) was determined. The mechanism of quinolone resistance in these isolates is being determined at the Antimicrobial Agents Research Group.

CLOSTRIDIUM DIFFICILE TOXIN B, BUT NOT TOXIN A, BINDS NUCLEIC ACID

S. Stewart and S. P. Borriello

Microbial Pathogenicity Group, Clinical Research Centre, Watford Road, Harrow, Middlesex, HA1 3UJ

In our studies on functions and mechanisms of action of *Clostridium difficile* toxins A and B we have looked at the interaction of these toxins with nucleic acid. Analysis of purified toxins A and B demonstrated that DNA co-purifies with toxin B following the final purification step by Mono Q anion exchange chromatography (MQEC), but that no nucleic acid was present in purified toxin A. The addition of protease (Pronase E; Sigma) to toxin B and analysis of the denatured product by MQEC showed that the DNA in the preparation no longer eluted where toxin B would. This finding showed that the point of elution of the DNA was dictated by intact toxin B and implied that this DNA was bound to toxin B as opposed to being unbound and simply co-eluting. The effect of removal of DNA from purified toxin B on the point of elution of toxin B has yet to be determined. However its effect on the cytotoxic activity has been investigated. The addition of DNAase (Sigma D-0876) consistently reduced the cytopathic effect of the toxin eight-fold, but had no effect on the cytotoxicity of toxin A. Use of a more purified DNAase (Sigma D-4527) still caused a reduction in toxin B cytotoxicity. The implications are that toxin B but not toxin A binds nucleic acid, and that its removal reduces the cytotoxic activity. The possibility that toxin B may bind target cell DNA, and that the feature of DNA binding may be shared by the related *C. sordellii* toxin LT, is being investigated.

MEASLES VIRUS AND CHRONIC ACTIVE HEPATITIS

J. D. Hayden, J. Teasdale, D. P. Jackson, J. Hoyle, J. I. Wyatt*, F. A. Lewis, D. I. Robertson,† H. Millward-Sadler,‡ M. F. Dixon and P. Quirke

*Department of Pathology, University of Leeds, *St James Hospital, Leeds, and Departments of †Pathology and ‡Medicine, University of Southampton*

Persistent measles virus infection causes subacute sclerosing panencephalitis (SSPE) and has been implicated in a range of chronic human diseases. Using the polymerase chain reaction (PCR), we have investigated its presence in SSPE and in chronic active hepatitis where a causal role has been suggested. Nucleic acid was extracted from paraffin-embedded tissue by standard techniques. With strict adherence to rigid anti-contamination procedures, nested reverse transcriptase PCR could reliably detect a 312-bp measles virus nucleocapsid RNA sequence in liver biopsies from two out of ten cases of chronic active hepatitis and in one well characterised case of autoimmune chronic active hepatitis. Under identical reaction conditions, the presence of measles virus nucleocapsid RNA was confirmed in two cases of wild type measles virus infection and in brain tissue from six cases of SSPE. Positive signals were absent from ten normal liver biopsies, ten cases of alcoholic hepatitis, ten cases of primary biliary cirrhosis and negative reagent controls containing distilled water in place of a nucleic acid

sample. High resolution polyacrylamide gel electrophoresis of products from successful amplifications showed them to be identical in length to fragments amplified from cloned measles virus nucleocapsid cDNA using the same primers. Direct sequencing and comparison of reaction products from a paraffin block of SSPE and cloned cDNA confirmed the amplification product to be measles virus. These results suggest a possible role for measles virus in chronic active hepatitis and will enable the detection and characterisation of RNA viruses in other chronic human diseases.

EXPRESSION OF EPSTEIN-BARR VIRUS TRANSFORMATION-ASSOCIATED GENES IN TUMOUR CELLS OF HODGKIN'S DISEASE

S. J. Hamilton-Dutoit, G. Pallesen, M. Rowe* and L. S. Young*

*Laboratory of Immunohistology, University Institute of Pathology, Finsensgade 12, DK-8000 Aarhus C, Denmark and *Department of Cancer Studies, University of Birmingham, Birmingham, B15 2TJ*

Recent DNA hybridisation studies have demonstrated Epstein-Barr virus (EBV) genomes in a substantial proportion of cases of Hodgkin's disease (HD), but it remains unclear whether the virus is involved directly in the pathogenesis of this tumour or is present merely as a passenger. The EBV-encoded latent gene products, latent membrane protein (LMP) and EBV nuclear antigen 2 (EBNA 2), play central roles in EBV-induced cell transformation *in vitro*, and have been implicated as important effector molecules in EBV-associated lymphomagenesis *in vivo*. We used immunohistochemistry with specific monoclonal antibodies to investigate the expression of these proteins in 84 cases of HD. LMP, but not EBNA 2, was demonstrated in Reed-Sternberg (RS) cells of 40 cases (48%). LMP expression varied with histological subtype, positive RS cells being seen in 1 (10%) of 10 cases of lymphocyte predominance, 16 (32%) of 50 cases of nodular sclerosis, and as many as 23 (96%) of 24 cases of mixed cellularity type. Our study (1) confirms the frequent association of EBV with RS cells in HD; (2) provides strong evidence of a direct oncogenic role for EBV in HD; and (3) indicates that LMP expression is related to histologically aggressive subtypes of HD. If future studies confirm an aetiological role for EBV, vaccination against the virus should be considered for the primary prevention of this malignancy.

CLONAL EBV GENOMES ARE DETECTABLE IN BIOPSIES FROM THE MAJORITY OF OLDER AND PAEDIATRIC HODGKIN'S DISEASE PATIENTS

Ruth Jarrett, D. Jones,* Alice Gallagher, Freda Alexander,† A. Krajewski,‡ Sarah Gledhill and D. Onions

*LRF Virus Centre, Department of Veterinary Pathology, University of Glasgow, *Department of Pathology, Southampton University Medical School, †LRF Centre for Clinical Epidemiology, University of Leeds, and ‡Department of Pathology, University of Edinburgh Medical School*

There is accumulating evidence that EBV is involved in the pathogenesis of a proportion of cases of Hodgkin's disease (HD). In this study we analysed the association between the presence of EBV genomes in tumour biopsies and the HD subtype and age of the patients. Of non-selected HD biopsies, 36% contained EBV genomes detectable by Southern blot analysis. Over 70% of biopsies from patients aged ≥ 50 years were EBV +ve ($n=38$) and over 50% of tumours from paediatric patients were positive ($n=13$). This contrasted with the results in the young adult group (aged 15–34) in which only 14% were EBV genome positive

($n=28$). These results show highly significant evidence for differences in EBV positivity by age. There was a slight excess of EBV positivity in the mixed cellularity HD cases as compared to the nodular sclerosing cases but this was not statistically significant. The detection of clonal EBV genomes in the majority of HD tumours from older and paediatric cases strengthens the argument that EBV is involved in the pathogenesis of these cases.

ANALYSIS OF EPSTEIN-BARR VIRUS (EBV) GENES AND GENE EXPRESSION IN AIDS-RELATED LYMPHOMA

S. J. Hamilton-Dutoit, G. Pallesen, K. Sandvej and L. S. Young*

*Laboratory of Immunohistology, University Institute of Pathology, Finsensgade 12, DK-8000 Aarhus C, Denmark and *Department of Cancer Studies, University of Birmingham, Birmingham, B15 2TJ*

HIV-infected individuals are at increased risk of developing lymphoma, some of which are associated with EBV. In-vitro EBV-infected cells may express several latent infection proteins (of which EBNA 2 and LMP show oncogenic transforming activity), a switch protein (ZEBRA) controlling induction of replication, and various replicative infection antigens (EA, VCA, MA). We studied the pathogenetic role of EBV in AIDS-related lymphoma (ARL) by investigating the presence of EBV-DNA (by in-situ hybridisation) and the pattern of EBV gene expression (by immunohistochemistry) in these tumours. We found: (1) EBV-DNA in 27 (43%) of 63 cases, viral genomes being found in 73% of immunoblast-rich ARL but in only 30% of Burkitt-type ARL; (2) EBV transformation-associated latent gene expression (LMP + EBNA 2) in 6 of 7 of EBV-DNA + lymphomas (but not in the single case of EBV-DNA + Burkitt-type ARL); (3) induction of ZEBRA in 5 of 7, EA in 3 of 7, and VCA and MA in 1 of 7 EBV-DNA + ARLs indicating replicative infection. Our results suggest that two groups of ARL exist: (a) immunoblast-rich tumours resembling EBV induced post-organ transplant lymphomas in morphology and viral gene expression, and in which EBV would appear to play a direct oncogenic role (contrary to most previous reports, EBV replicative infection may occur in this group suggesting that anti-viral therapy may be appropriate); (b) Burkitt-type, associated with EBV in only a minority of cases in which other oncogenic factors appear to be of greater importance.

MYCOBACTERIUM PARATUBERCULOSIS IN INFLAMMATORY BOWEL DISEASE

P. Quirke, D. Dockey, P. Hawkey,* F. A. Lewis, D. Graham,† G. R. Taylor‡ and M. F. Dixon

*Departments of Pathology and *Microbiology, University of Leeds, †Baylor College of Medicine, Houston, Texas, USA and ‡Yorkshire Regional DNA Laboratory, Leeds General Infirmary*

Mycobacterium paratuberculosis causes Johne's disease in cattle and has been implicated in the causation of human Crohn's disease. Using nested PCR we have designed primers to the IS900 insertional element and the gene encoding 16S RNA of *M. paratuberculosis*. These primers have been tested on specific *M. paratuberculosis* isolates from cases of Crohn's disease, Ben and Dominic, together with *M. avium intracellulare* and *M. tuberculosis*. Using IS900 3' and 5' sets of primer pairs we have amplified *M. paratuberculosis*, Ben and Dominic, but not *M. avium intracellulare* or

M. tuberculosis. 16S RNA primers have identical 16S RNA sequences to *M. avium intracellulare* and, therefore, also amplified this organism but not *M. tuberculosis*. DNA extracted from paraffin embedded material from five cases

of Johne's disease of cattle were successfully amplified by all three sets of primers, confirming the presence of *M. paratuberculosis*. Early results on human Crohn's disease are encouraging.

POSTER PRESENTATIONS

DEMANDS OF AN EPIDEMIC OF RESPIRATORY SYNCYTIAL VIRUS (RSV) INFECTION ON HOSPITAL SERVICES

B. I. Duerden

Microbiology Department, Children's Hospital, Western Bank, Sheffield

During Oct. 1989–Feb. 1990, RSV bronchiolitis was confirmed by immunofluorescent antibody tests (IFAT) on nasopharyngeal aspirates in 182 infants <1 y (mean 4.7 m) admitted to the Children's Hospital, Sheffield (pop. 500 000; cf 93 admissions in 1988–89). The epidemic peak was 30 cases in the second week of December when routine admissions were suspended. Mean hospitalisation was 4.8 d. Most (78%) were mild cases, recovering quickly (mean 3.8 d) with only nasogastric tube feeding (36%) or no active management; only 15% of these were given antibiotics. A further 13% were moderately severe and required O₂ by head-box (12.5%), i.v. fluids (7%) and nasogastric feeds; 60% of the moderate cases received antibiotics. Only 6% of cases were severe, requiring admission to the intensive care unit (ICU), O₂, ventilation (IPPV), i.v. fluids and i.v. antibiotics. There were five deaths—two in infants with congenital abnormalities and three presenting as sudden infant death syndrome (two with no histological evidence of bronchiolitis). Bacterial pathogens (*Haemophilus influenzae* 3, pneumococcus 2, *Branhamella catarrhalis* 6, *Klebsiella* sp. 2, *Staphylococcus aureus* 1) were isolated from eleven patients. Overall, 26% of patients were given antibiotics: amoxycillin in 19% (+ flucloxacillin in 9%), others in 7%. The epidemic represented 859 bed-days, 195 with close monitoring on wards and 70 in ICU. Rapid IFAT diagnosis reduced laboratory investigations, avoided general use of antibiotics and enabled early, confident discharge of most patients.

A STUDY OF HUMAN RESPIRATORY TRACT CHLAMYDIAL INFECTIONS IN CAMBRIDGESHIRE 1986–88

T. G. Wreghitt, C. E. Barker, J. D. Trehanne,* J. M. Phipps,* V. Robinson† and R. B. Buttery‡

*Clinical Microbiology and Public Health Laboratory, Addenbrooke's Hospital, Cambridge; *Section of Virology, Institute of Ophthalmology, Judd Street, London; †Department of Pathology, University of Cambridge; and ‡Public Health Medicine, Cambridge Health Authority*

Human respiratory tract chlamydial infections have been studied in Cambridgeshire for many years, but until recently we have been unable to distinguish between infection with *Chlamydia psittaci* or *Chlamydia pneumoniae*. In this study, we have employed the micro-immunofluorescence (micro-IF) test for this purpose and to look for the relative incidence of *C. psittaci* and *C. pneumoniae* infections in Cambridgeshire. Among 50 patients with community-acquired respiratory tract symptoms whose serum samples had chlamydia complement fixation test titres ≥ 64 , 25 had evidence of

recent *C. psittaci* or *C. pneumoniae* infection. Nineteen (76%) of the 25 patients had evidence of recent *C. psittaci* infection and of these 16 (84%) had recently had contact with birds. Six patients (24%) had evidence of recent *C. pneumoniae* infection, and of these, only two (33%) had recently had contact with birds. Whereas *C. psittaci* was isolated from several of the birds associated with human *C. psittaci* infection, it was not cultured from any of the birds in contact with the two human *C. pneumoniae* cases.

A RADIOCHEMICAL METHOD FOR MEASURING ADHESION OF COAGULASE-NEGATIVE STAPHYLOCOCCI (CNS) TO SURFACES OF PLASTIC OR GLASS

M. Hussain, J. G. M. Hastings* and P. J. White

*Department of Molecular Biology and Biotechnology, The University of Sheffield, P.O. Box 594, Firth Court, Western Bank, Sheffield, S10 2UH and *Department of Clinical Microbiology, Queen Elizabeth Hospital, Birmingham*

We have already reported that growth of CNS in a chemically defined medium (HHW) containing U-[¹⁴C] glucose leads to heavy labelling of the extracellular slime. A firmly adherent mass of slime plus organisms accumulates on the sides of the culture tubes when strains of CNS are grown in radioactive medium HHW. This mass was washed (to remove labelled medium) and then counted after adding scintillation fluid. Organisms from the liquid culture also were washed and counted to check that ¹⁴C-glucose had been utilised. Nine strains of CNS were examined in this way, and the results were compared with those of five older techniques for measuring adherence. The new method is quick, and has advantages of reproducibility and good discrimination between strains; there is a 15-fold difference in adherent counts between slime producing and non-producing strains. With the new assay, the effects of several antibacterial compounds were investigated, at levels below their minimum growth-inhibitory concentrations, on attachment of four slime-producing strains. Tunicamycin, chloramphenicol and 5-fluorouracil each totally prevented adherence while several other drugs had much less effect.

COMMON EPITOPES IDENTIFIED BY MONOCLONAL AND POLYCLONAL ANTIBODIES AGAINST 70-KDA IRON-REGULATED OUTER-MEMBRANE PROTEINS OF NEISSERIA MENINGITIDIS: THEIR SURFACE EXPOSURE AND FUNCTIONAL ROLE

D. A. Ala'Aldeen, Heather A. Davies and S. P. Borriello

Microbial Pathogenicity Research Group, MRC Clinical Research Centre, Watford Road, Harrow, HA1 3UJ

The 70-Kda iron regulated protein (FeRP-70) is the most prominent of the several new outer-membrane proteins expressed by *Neisseria meningitidis* when grown under iron limitation. We have purified FeRP-70 by elution from SDS-polyacrylamide gels and raised rabbit monospecific poly-

clonal and murine monoclonal antibodies against it. On Western blots these anti-FeRP-70 antibodies cross-react with common epitopes on FeRP-70's from other serogroups and serotypes of *N. meningitidis* and some commensal neisseria. Immunogold electronmicroscopy, with polyclonal and monoclonal antibodies and whole cells and sections, did not reveal increases in labelling, suggesting lack of surface exposure. None of these antibodies exhibited bactericidal activity. Although these common epitopes appear immunogenic *in vivo*, their lack of surface exposure and absence of activity by antibodies directed against them, do not support a potentially protective role for these antigens as a component of a vaccine.

RECOGNITION OF ISOLATES OF THE *CANDIDA ALBICANS* DARLINGTON STRAIN BY RFLP ANALYSIS

Michelle A. Pearce, S. A. Howell and W. C. Noble

Department of Microbial Diseases, Institute of Dermatology, St Thomas' Hospital, Lambeth Palace Road, London, SE1 7EH

The *Candida albicans* Darlington strain was the first reported clinically azole-resistant strain in the UK; subcultures were, therefore, passed between laboratories in studies to elicit the mechanism of resistance. Seven isolates representing subcultures of this strain were obtained from several laboratories, including a clinical *C. albicans* isolate from the patient in 1988 and one isolate that was known to have become azole sensitive. These proved to have a unique sterol pattern. RFLP patterns of total DNA from these isolates were compared and 5 of the 7 were found to be identical. The two isolates exhibiting different RFLP patterns included the sensitive variant and one other subculture; both were found to be azole sensitive and had different RFLP patterns from each other.

SDS-PAGE CHARACTERISATION OF SPECIES OF THE GENUS *PREVOTELLA*

R. Yousefi and B. I. Duerden

Department of Experimental and Clinical Microbiology, University of Sheffield Medical School, Beech Hill Road, Sheffield, S10 2RX

Species of the melaninogenicus-oralis group of mainly oral *Bacteroides* have now been separated into a new genus, *Prevotella*. Many are phenotypically similar and cannot be readily distinguished by routine tests. Twenty-one reference strains (*P. melaninogenica* NCTC 11139, *P. denticola* ATCC 35308, *P. ivescheii* ATCC 15930, *P. intermedia* NCTC 9338, *P. corporis* ATCC 33574, *P. oralis* NCTC 11459, *P. veroralis* ATCC 33779 and 83-457F, *P. buccalis* ATCC 35310, *P. pentosacea* NP 333 and WPH 61, *P. oris* ATCC 33573 and 86-201DD, *P. buccae* ATCC 33574 and 83-617, *P. zoogloeiformans* ATCC 33285, *P. bivia* VPI 5540, *P. disiens* NCTC 11157 and VPI 8057, *P. oulora* NCTC 11871 and *P. capilla* ATCC 33690) and 91 non-pigmented isolates from adult periodontal disease were examined by conventional biochemical tests, gas-liquid chromatography (GLC) and enzyme tests, and whole-cell protein profiles were obtained by sodium dodecyl sulphate-polyacrylamide gel electrophoresis (SDS-PAGE). There was close correlation between patterns of results in biochemical, GLC and enzyme tests and the SDS-PAGE profiles. The reference strains of the designated species gave distinct SDS-PAGE profiles. The 91 periodontal isolates were assigned to nine groups by their biochemical test patterns and SDS-PAGE profiles; the profiles of most (79) isolates were very similar to those of

seven species reference strains (*P. buccae* 42, *P. veroralis* 15, *P. oralis* 8, *P. oris* 7, *P. bivia* 4, *P. buccalis* 1, *P. zoogloeiformans* 1) and 13 isolates remained in two undesigned SDS-PAGE groups. Minor variations in profiles enabled the recognition of sub-groups within some species.

SPONGIFORM ENCEPHALOPATHY: CAN IT BE CURED?

S. F. Dealler

Microbiology Department, University of Leeds, Old Medical School, Leeds, LS2 9JT

Transmissible spongiform encephalopathy (TSE) involves the presence in infected brain and other tissues of an ultra-filterable agent, which is resistant to heat, ultraviolet light, ionising radiation, chemical disinfectants, DNAase, RNAase, proteinases, and chemicals specifically active against nucleic acid. The finding of a protein (PrP) in infected tissue that is not present in non-infected tissue and which is associated with infective structures (scrapie associated fibrils) has suggested to researchers that this is the infective agent (prion). However, few adequate explanations as to how this agent can cause its own multiplication have been put forward. This presentation explained the 'crystal' hypothesis and how this would allow potential treatment of the disease. Crystalline forms of polyPrP appear in neurone and microglia as mats of parallel electron-dense fibres. As other crystals, they would be expected to allow further molecules to join only if their molecular pattern specifically fitted the crystal. A method of preventing crystal extension using chemicals was explained.

BREAKDOWN OF β -LACTAM COMPOUNDS BY β -LACTAMASES FROM SENSITIVE *ESCHERICHIA COLI* STRAINS

R. Edwards, F. Soriano* and D. Greenwood

*Department of Microbiology and PHLS Laboratory, University Hospital, Queen's Medical Centre, Nottingham, NG7 2UH and *Department de Microbiologia, Fundacion Jimenez Diaz, Avda. Reyes Catolicos 2, 28040 Madrid, Spain*

In common with other gram-negative bacilli, virtually all *Escherichia coli* strains possess a chromosomally-mediated β -lactamase which is able slowly to hydrolyse some early cephalosporins, such as cephalothin and cephaloridine. It has been assumed that later β -lactam compounds that are highly stable to most plasmid-mediated β -lactamases would not be hydrolysed by this chromosomal enzyme. In an investigation of inoculum effects with ceftizoxime and some other β -lactam agents, it was noticed that ceftizoxime activity was abolished during overnight incubation with high inocula of ostensibly sensitive *E. coli*. This prompted us to investigate the comparative stability of some representative β -lactam agents to crude extracts of sensitive strains of *E. coli*. Two strains were tested: *E. coli* ATCC 25922 and a fresh clinical isolate, *E. coli* S31. Aqueous solutions (100 mg/L) of ampicillin, cefuroxime, ceftizoxime, cefminox (a new cephamycin), aztreonam and imipenem were incubated at 37°C in a 1:1 ratio with crude sonicated extracts of each strain. Samples were removed at intervals for assay by high pressure liquid chromatography. Ceftizoxime was completely hydrolysed by crude extracts of both strains within 24 h; less than 20% of ampicillin or cefminox remained after 24 h. The effect on cefuroxime and aztreonam appeared to be variable; neither strain was able to break down imipenem. These results show that crude β -lactamase extracts from β -lactam sensitive *E. coli* strains have the capacity to break down certain β -lactam antibiotics when tested at high concentrations *in vitro*.

THE EFFECT OF ANAEROBIOSIS ON THE ACTIVITY OF GENTAMICIN AND FOSFOMYCIN AGAINST *ESCHERICHIA COLI*

J. Brown, R. Edwards and D. Greenwood

Department of Microbiology and PHLS Laboratory, University Hospital, Queen's Medical Centre, Nottingham, NG7 2UH

The effect of anaerobiosis on the activity of antibiotics may be of clinical importance since many infected sites exist under conditions of reduced oxygen potential. Anaerobic conditions have long been known to impair the activity of aminoglycoside antibiotics by interfering with the uptake of these drugs into target cells. In contrast, it has been reported that anaerobiosis can potentiate the antibacterial effect of fosfomycin, a drug known to display better activity *in vivo* than tests *in vitro* would indicate. In the present study, continuous turbidimetric monitoring and conventional minimum inhibitory concentration (MIC) techniques were used to investigate the influence of anaerobic conditions on the activities of fosfomycin and the aminoglycoside antibiotic, gentamicin. Standard agar incorporation titration confirmed previous findings in showing MICs of gentamicin to be raised and MICs of fosfomycin to be lowered for a variety of bacteria tested under anaerobic conditions. However, turbidimetric experiments with two clinical isolates of *E. coli*, in which a considerably higher bacterial inoculum was used (10^8 cfu/ml compared to 10^6 cfu/spot for the agar incorporation MIC method) showed no significant difference between the antibacterial effects of fosfomycin under the two atmospheric conditions. The activity of gentamicin on dense bacterial populations was reduced about twofold in tests carried out under anaerobic conditions, a smaller reduction in activity than was observed using agar incorporation MIC titration.

HERPES SIMPLEX VIRUS TYPE I IN HUMAN DONOR CORNEAS

R. E. Bonshek, B. Marcyniuk,* C. Dennett,† A. B. Tullo,* P. E. Klapper,† G. M. Cleator† and A. G. Lewis†

*Divisions of Molecular Pathology and †Virology, Department of Pathological Sciences, Manchester University, Oxford Road, Manchester, M13 9PT and *Royal Eye Hospital, Oxford Road, Manchester, M13 9WH*

Corneal transplantation is an accepted ophthalmological procedure with a low incidence of graft failure due to rejection or other complications. Whilst the potential for the transmission of certain infectious agents exists, this is seen only very rarely. A number of agents has been reported to have been transmitted by this procedure, but there is no conclusive evidence that Herpes Simplex Virus type 1 (HSV-1), a common ocular pathogen, has been transmitted in this way. In the present study, HSV-1 was identified in the stromal cells of stored human corneal donor material. In both corneas from one donor and in one cornea from a second donor, complete and unaccountable endothelial cell loss occurred. Culture of tissue storage medium was undertaken and HSV-1 was isolated. Examination of fixed corneas by immunocytochemistry, transmission electronmicroscopy and in-situ DNA hybridisation revealed viral antigen and virus particles, and viral DNA hybridisation revealed viral antigen, virus particles and viral DNA (respectively) in keratocytes. These findings are of significance for corneal transplantation and serious consideration must be given to the possibility of transmission of HSV-1 by this procedure, particularly in situations where fresh donor tissue is used or where corneas have been kept in short-term storage.