Isolation and purification of *Aeromonas sobria* cytotoxic enterotoxin and β -haemolysin

P. J. GOSLING, P. C. B. TURNBULL*, N. F. LIGHTFOOT†, J. V. S. PETHER and R. J. LEWIS‡

Public Health Laboratory, Musgrove Park Hospital, Taunton, Somerset TA1 5DB, *PHLS Centre for Applied Microbiology and Research, Division of Biologics, Porton Down, Salisbury, Wiltshire SP4 0JG, †Public Health Laboratory, General Hospital, Westgate Road, Newcastle upon Tyne NE4 6BE and ‡Faculty of Applied Sciences, University of the West of England, Frenchay, Bristol BS16 1QY

Summary. *Aeromonas* sp., grown in tryptone soya broth supplemented with yeast extract, 0.6%, pH 7.5, and incubated with agitation at 100 oscillations/min for 15 h at 37°C produced optimal amounts of β -haemolysin and cytotoxic enterotoxin. More prolonged incubation resulted in the loss of enterotoxic activity and anion exchange chromatographic analysis indicated the presence of a moiety capable of breaking down the toxin. Anion exchange fast protein liquid chromatography resulted in a single peak of haemolytic activity and two peaks with enterotoxic activity. The cytotoxic enterotoxin was purified from the fraction most active in the infant mouse assay; the second peak, which did not cross-react immunologically, may represent a second cytotoxic enterotoxin. Neither peak was observed in the chromatographic fractions of filtrates from strains devoid of activity in the infant mouse assay. Purified enterotoxin, estimated to have a mol. wt of 15 kDa by SDS-PAGE, caused fluid accumulation in the infant mouse assay, was non-haemolytic to rabbit erythrocytes, caused an increase in cAMP activity in tissue culture cells and did not cross-react immunologically with components of cholera toxin or the whole toxin. Purified β -haemolysin had an estimated mol. wt of 55 kDa, lysed rabbit erythrocytes and did not cause fluid accumulation in the infant mouse test.

Introduction

Motile *Aeromonas* spp. have been incriminated as aetiological agents of diarrhoeal disease in man for two decades.^{1–4} Isolates of motile aeromonads from human faeces have been associated significantly with symptoms of gastroenteritis,⁵ shown to produce extracellular products that elicit an enterotoxic response in the infant mouse assay⁶ and to produce adhesins to human intestinal cells.⁷ However, attempts to fulfil Koch's postulates by reproducing the disease in animal models, or even in man, have been unsuccessful.⁸

To evaluate the potential of *Aeromonas* spp. as enteropathogens, a better understanding of their pathogenesis is necessary. Progress has been hampered by confusion in the taxonomy of the genus and by the complex extracellular protein profile of members of the genus. The biologically active proteins produced by motile aeromonads include haemolysins and at least one enterotoxin that are considered to be potential virulence factors in *Aeromonas*-associated diarrhoeal disease.

In previous studies on *Aeromonas* enterotoxin,⁸ crude preparations have been used and attempts to assess their role as virulence factors have been com-

plicated by the presence of other extracellular products. It remains unclear whether there is one or more *Aeromonas* enterotoxins, whether the toxin is cytotoxic, i.e. causing tissue damage, cytotoxic, i.e. exerting an enterotoxic effect without tissue damage, or if it cross-reacts immunologically with cholera toxin.

The aims of the present study were to establish the optimum conditions for enterotoxin production, and hence detection of enterotoxigenic strains, and to separate and purify aeromonas enterotoxin and haemolysin to clarify their roles as virulence factors in disease in man.

Materials and methods

Bacterial strains

Two strains of *Aeromonas* spp. (VL8509 and VL8804) shown previously to produce haemolysin and enterotoxin,⁹ and two strains (VL7711 and VL8778) shown to be devoid of these activities, were kindly supplied by Dr J. V. Lee, PHLS Centre for Applied Microbiology and Research, Porton Down, Wiltshire. Strains were maintained on Nutrient Agar slopes overlaid with sterile oil at room temperature and in

Nutrient Broth (Unipath) containing glycerol 20% at -70°C .

Preparation of cell-free culture filtrates (CFCF)

Fresh subcultures of *Aeromonas* strains on nutrient agar were used to inoculate 10 ml of tryptone soya broth (Unipath) supplemented with yeast extract 0.6% in 100-ml conical flasks which were incubated in a flask shaker at various times and temperatures. Cultures were then centrifuged at 10000 *g* for 10 min at 4°C and the supernate was filtered through a 0.22- μm low protein binding membrane filter (Sartorius GmbH, Germany). CFCF were stored at -70°C . To establish parameters during toxin production that could affect detection of enterotoxigenic strains, experiments varying the pH, agitation rate, time and temperature of growth were conducted. Growth curves were constructed from viable counts obtained by a standardised drop count method¹⁰ at timed intervals.

Concentration of CFCF

To facilitate detection of enterotoxic activity the CFCF were concentrated 20-fold by ultrafiltration through an Amicon PM series filter apparatus (Amicon, Basingstoke) with a Diaflo membrane PM10 having a 10-kDa cut off. Ultrafiltration was performed in an ice bath.

Enterotoxin assay

Enterotoxic activity was determined in infant mice. The mice were inoculated by intraperitoneal injection with 0.1 ml of CFCF to which Evans blue 2% solution had been added (10 μl in 1 ml of CFCF). Each test involved two to five mice and the intestinal weight/body weight (IW/BW) ratio was determined at $3\text{ h} \pm 10\text{ min}$. An IW/BW ratio of > 0.8 was considered to indicate enterotoxic activity.

Haemolysin assay

Doubling dilutions of unconcentrated CFCF in normal saline, or of fractions following FPLC separations, were made and 100- μl volumes were added to equal volumes of a suspension of rabbit erythrocytes 0.8% in normal saline in 96-well round-bottomed trays. Haemolysis was recorded after incubation for 1 h at 37°C and then after subsequent incubation for 2 h at 4°C . The highest dilution giving complete haemolysis was taken as the end point.

Protein determination

Protein was measured by the Folin-Lowry method and with a protein assay kit (BioRad) with bovine serum albumin as a standard.

Chromatography methods

A fast protein liquid chromatography (FPLC, Pharmacia, Milton Keynes) apparatus was used.

Anion exchange chromatography was performed on a pre-packed Mono Q column HR 5/5 (Pharmacia). Buffer A consisted of 30 mM triethanolamine buffer, pH 8, and buffer B of 30 mM triethanolamine buffer, pH 8, containing 1 M NaCl. All reagents were of analytical grade and all buffers and solutions were pre-filtered through a 0.22- μm filter before use. Various elution programmes were constructed to vary the linear gradient rate from 0 to 100% buffer B with a flow rate of 1 ml/min. The protein content of the eluate was monitored at 280 nm and all fractions over the course of the gradient were collected.

Gel filtration was performed on a pre-packed column HR 10/30 (Pharmacia) containing 24 ml of Superose 12. The elution buffer consisted of 0.05 M NaH_2PO_4 , pH 7.0, containing 0.15 M NaCl and the flow rate was 0.5 ml/min. The protein content of the eluate was monitored at 280 nm and all fractions were collected throughout the separation. Immediately after separation, with the same elution buffer, mol. wt standards were applied to the column and the elution times were noted. The standards consisted of bovine serum albumin (67 kDa), ovalbumin (43 kDa), chymotrypsinogen (25 kDa) and ribonuclease (13.7 kDa) (Sigma).

*Scale-up of production and purification of *Aeromonas* toxins*

Batches of CFCF (250 ml) were pooled until 2 L was available. This was diluted four-fold with 30 mM triethanolamine buffer containing 1 mM EDTA, pH 8.0 (buffer C), and an anion exchange resin (Q Sepharose Fast Flow, Pharmacia) was added and kept suspended for 1 h at 4°C . This was then allowed to settle at 4°C and the resin was collected in a large diameter chromatography column. The resin was then washed with three bed volumes of pre-cooled buffer C. Protein was eluted from the resin with three bed volumes of buffer C containing 0.23 M NaCl at 4°C . Three sequential protein elutions from the resin were performed by washing with three bed volumes of buffer C containing 0.23 M, 0.48 M and 1 M NaCl respectively. The products were collected and stored at -70°C . Subsequently they were subjected separately to precipitation with saturated ammonium sulphate 70% and the precipitate was collected by centrifugation at 23000 *g*. This was re-suspended in 2 ml of buffer C and dialysed at 4°C for 5 h against a large volume of buffer C. Insoluble material was removed by centrifugation at 15000 *g* and the fluids were filter sterilised through a 0.22- μm filter. The preparation originating from the wash-through with 0.23 M NaCl was regarded as crude aeromonas β -haemolysin and that originating from the wash-through with 1 M NaCl as crude aeromonas enterotoxin. The crude toxin

Fig. 1. Haemolytic (▨) and enterotoxin (▩) activity in CFCF of *A. sobria* strain VL8509 sampled during growth in tryptone soya broth supplemented with yeast extract 0.6%.

preparations were purified by application to a pre-packed Mono Q HR 16/10 (Pharmacia) anion exchange column and active fractions were collected. Buffer exchange was carried out by ultrafiltration with filters of 10000-kDa cut off (Sartorius) and the active fractions were again subjected to anion exchange chromatography as before. Active fractions were collected, dialysed against phosphate buffered saline (PBS), pH 7.4, and the resulting purified toxins were stored at -70°C .

Gel electrophoresis

SDS-PAGE was performed as described by Laemmli¹¹ on CFCF samples and purified toxins. Samples in SDS reducing buffer (0.5 M Tris-HCl, pH 6.8, containing SDS 2% w/v, glycerol 10% v/v, 2-mercaptoethanol 5% v/v and a trace of bromophenol blue dye) were boiled for 2 min and applied to a 1.5-mm resolving gel of acrylamide 12% w/v in 375 mM Tris-HCl, pH 8.8, containing SDS 0.1% *via* a stacking gel, used to concentrate samples to improve band resolution, consisting of acrylamide 4% w/v in 125 mM Tris-HCl, pH 6.8, containing SDS 0.1% w/v. Rabbit muscle phosphorylase b (106 kDa), bovine serum albumin (80 kDa), hen egg white ovalbumin (49.5 kDa), bovine carbonic anhydrase (32.5 kDa), soya bean trypsin inhibitor (27.5 kDa) and hen egg white lysozyme (18.5 kDa) were included as mol. wt markers (BioRad).

Electrophoresis with a voltage gradient of 80 V was performed in 25 mM Tris-HCl (pH 8.3) containing 192 mM glycine and SDS 0.1% w/v. Gels were stained overnight with Coomassie Brilliant Blue R250 0.1%

w/v in methanol: acetic acid:water (40:10:50) and cleared in several changes of the same solvent.

Ganglioside GM1 binding assay

Ganglioside GM1 (Sigma) 50 $\mu\text{g/ml}$ diluted in carbonate buffer, pH 9.6, was added to each well of a 96-well round-bottomed tray. After overnight incubation at 4°C , the trays were washed three times with 0.1 M PBS containing Tween 20 0.5% (PBST), pH 7.4, and bovine serum albumin 0.1%. The trays were washed once with distilled water and dried. Test solutions (50 μl) of crude or purified aeromonas toxins were added to wells and incubated at room temperature overnight. Solutions containing cholera toxin (CT; Sigma), 0.5–500 mg/ml, and uninoculated broth were also included as positive and negative controls, respectively. All solutions were tested in duplicate. After three washes with PBST, goat anti-CT (Calbiochem, Nottingham) and rabbit anti-aeromonas cytotoxic enterotoxin at a dilution of 1 in 640 were used to detect bound antigen. Diluted normal rabbit serum, 1 in 10–1 in 1280, was added to duplicate wells treated with test samples to provide negative controls. After overnight incubation at 4°C , the trays were washed three times with PBST and protein A peroxidase conjugate (Sigma) diluted 1 in 1000 was added to detect bound immunoglobulins.

After incubation for 1 h at 37°C , 100 μl of substrate, 2,2'-azino-di-(3-ethylbenzthiazoline sulphonic acid) (ABTS) was added to each well. Trays were incubated for 30 min at 37°C and the absorbance of wells was read at $A_{410\text{ nm}}$ with a MicroELISA Minireader (Dynatech). Wells were considered to have demon-

Fig. 2. The effect of (a) duration of incubation, (b) agitation, (c) temperature of incubation and (d) starting pH, of 15-h cultures of *A. sobria* strain VL8509 in tryptone soya broth supplemented with yeast extract 2% on the elaboration of enterotoxin. Each result represents a mean value obtained from three mice and vertical lines indicate SD. Intestinal weight/body weight (IW/BW) ratios above the dotted line are considered an enterotoxigenic response.

strated binding to ganglioside GM1 at an A_{410} reading of > 0.1 .

Cyclic AMP determination

Stimulation of cAMP production in Chinese hamster ovary tissue culture cells treated with crude and purified aeromonas cytotoxic enterotoxin was determined. Tissue culture cells were grown for 24 h in 25-ml flasks, treated with 0.33 ml of the toxin and incubated at 37°C overnight. Three flasks were used for each sample and the cells were trypsinised, pooled for each sample, rinsed in PBS, pH 7, and homogenised in 4 mM EDTA, pH 7. Samples were tested in duplicate. The sonicates were boiled for 3 min, centrifuged for 5 min and the amount of cAMP in 50 µl of supernate was determined by radio-immunoassay, kindly carried out by Dr P. J. Wood at the Regional Endocrine Laboratory, Southampton. Cholera toxin at 5 and 50 ng/ml, untreated cells and 20-fold broth

concentrates were used as positive and negative controls, respectively.

Stability of toxins

Purified toxin (in 0.2-ml volumes) was incubated with trypsin, pronase and papain (Calbiochem, 1.0 mg/ml) for 15 min at room temperature and then tested for activity. Sample volumes were exposed to a range of temperatures (25°–80°C) in glass tubes in a waterbath for 15 or 60 min and cooled rapidly before testing for remaining haemolytic and enterotoxic activity.

Immunological cross-reactivity test

Gel immunodiffusion tests were performed in agarose gel (BDH Chemicals Ltd) 1% w/v. Purified toxins were tested against specific antisera and antisera to whole cholera toxin, cholera A subunit and cholera B subunit (Calbiochem) in 3 mm diameter wells.

Antisera were tested neat and at five- and 50-fold dilutions in PBS, pH 7.4.

Results

Production of toxins

The pH, agitation rate, time and temperature of growth giving optimal toxin yields by strains VL8509 and VL8804 were determined (figs. 1 and 2). Optimum enterotoxigenicity was observed after growth in tryptone soya broth supplemented with yeast extract 0.6%, pH 7.5, for 15 h at 37°C with agitation of 100 oscillations/min. Enterotoxigenicity became detectable in the

Table I. Mean haemolytic and enterotoxigenic activity of 20 human faecal isolates after incubation for 15 and 17 h

Incubation time (h)	Haemolytic activity —mean(SD)HU/ml	Enterotoxigenic activity —mean(SD)IW/BW ratio
15	$8.2 (3.1) \times 10^2$	$8.9 (90.4) \times 10^2$
17	$9.2 (3.3) \times 10^2$	$6.8 (0.6) \times 10^2$

HU, one haemolytic unit is the dilution that caused complete haemolysis of rabbit erythrocytes.

IW/BW, intestinal weight/body weight ratio.

Fig. 3. FPLC gel filtration separation of an enterotoxigenic CFCF of *A. sobria* strain VL8509: haemolytic (▨) and enterotoxigenic (▩) activity.

Table II. Purification of cytotoxic enterotoxin from *A. sobria* strain VL8509

Method of purification	Volume (ml)	Protein content (mg/ml)	Enterotoxigenic activity (EU/ml)	Specific activity (EU/mg)	Total activity (EU $\times 10^3$)	Purification ratio	Yield (%)
Cell-free culture filtrate	2000	4.2	40	9.5	3360	1.0	100
Sephadex Fast Flow (crude toxin)	33	5.5	1000	181.8	330	19.1	9.8
FPLC Mono Q anion exchange chromatography (purified toxin)	4.2	1.2	500	416.7	21	43.9	0.6

EU, one enterotoxigenic unit is the dilution that caused an IW/BW ratio of > 0.08 in the infant mouse assay.

Fig. 4. FPLC anion exchange chromatograph of an enterotoxigenic CFCF of *A. sobria* strain VL8509: haemolytic (▨) and enterotoxigenic (▩) activity.

CFCF during the late logarithmic phase of growth after incubation for 12 h, reaching a peak at 15 h. However, this activity disappeared after 17 h. Haemolytic activity became detectable after 5 h and remained throughout the experiment, reaching a peak at 19 h.

Confirmation of these results was obtained in tests on the CFCF of 20 human faecal isolates of *Aeromonas* spp. for haemolytic and enterotoxigenic activity after incubation for 15 and 17 h (table I). Enterotoxin was not evident at 12–17 h and haemolysin was not detected from 5 h onwards in the CFCF of strains VL7711 and VL8778, which had been included as negative controls on the basis of previous findings.⁹

Anion exchange chromatography of CFCF revealed changing protein profiles with prolonged incubation and subsequent loss of enterotoxigenicity. The lack of haemolytic and enterotoxigenic activity by strains VL7711 and VL8778 was confirmed under all the conditions tested.

Gel filtration

A large band of proteins was eluted from the Superose 12 HR 10/30 column with 12–20 ml of elution buffer. Haemolytic activity was observed over a band of eight fractions eluted with 10–18 ml of buffer; enterotoxigenic activity was detected in a single fraction eluted with 17 ml of buffer. The enterotoxigenic

Table III. Purification of haemolysin from *A. sobria* strain VL8509

Method of purification	Volume (ml)	Protein content (mg/ml)	Haemolytic activity (HU/ml)	Specific activity (HU/mg)	Total activity (HU × 10 ⁴)	Purification ratio	Yield (%)
Cell-free culture filtrate	2000	4.2	1024	244	204.9	1.0	100
Sephadex Fast Flow (crude toxin)	24	6.5	16384	2520.6	39.3	10.3	19.2
FPLC Mono Q anion exchange chromatography (purified toxin)	3.2	2.2	4096	13107.2	9.2	53.7	4.5

Table IV. Susceptibility of aeromonas toxins to temperature and enzymes

Temperature (°C)	Time (min)	Activity remaining (%)		Other treatment (15 min)	Activity remaining (%)	
		Haemolysin	Enterotoxin		Haemolysin	Enterotoxin
25	15	100	95	Trypsin 1 mg/ml Pronase 1 mg/ml Papain 1 mg/ml	100	96
25	60	95	92			
35	15	95	82		95	90
35	60	75	60			
45	15	85	42		5	0
45	60	20	27			
55	15	5	0			
55	60	0	0			
65	15	0	0

Fig. 5. SDS-PAGE of extracellular proteins of *A. sobria* strain VL8509 stained with Coomassie Blue. (a) Lane 1, purified cytotonic enterotoxin; 2, crude CFCF; 3, mol. wt markers. (b) Lanes 1 and 3, purified haemolysin; 2, mol. wt markers.

fraction was not distinct from the haemolytic fraction (fig. 3) when separation was attempted by this method.

Anion-exchange chromatography

Assay of each individual fraction eluting from

the Mono Q HR 5/5 FPLC column for haemolytic and enterotoxic activity showed that the aeromonas β -haemolysin that caused complete haemolysis of erythrocytes was contained in a single fraction eluted with 0.17 M NaCl (fig. 4). Partial haemolysis was also detected over two broad bands. The first band eluted

Table V. cAMP production by CHO tissue culture cells treated with various toxin preparations

Treatment	cAMP activity (pmol/ml)	Increase in activity (fold)
Tryptone soya broth with 0.6% yeast extract (control)	1.1	1.0
Cholera toxin 50 ng	13.7	11.5
Crude aeromonas enterotoxin	7.5	5.8
Purified aeromonas enterotoxin	8.5	6.7
Cell-free culture filtrate of strain VL8509	0.9	0

over five fractions with 0.27–0.38 M NaCl, with the peak of activity at 0.34 M NaCl. The second broad band occurred over four fractions eluting with 0.42–0.51 M NaCl, with the peak of activity at 0.44 M NaCl.

Two single fractions eluting with 0.55 M NaCl and 0.85 M NaCl, respectively, caused fluid accumulation in the infant mouse test (fig. 4). The enterotoxic fractions were not haemolytic and the haemolytic fractions were not enterotoxic. The increase in specific activity and yields during toxin purification can be seen in tables II and III.

Characteristics of purified toxins

Aeromonas cytotoxic enterotoxin and aeromonas β -haemolysin appeared as single bands in SDS-PAGE gels (fig. 5) with corresponding mol. wts of 15 and 55 kDa respectively. Both were heat-labile and were inactivated at 55°C for 15 min and by papain but not by trypsin or pronase (table IV). Purified toxins did not cross-react immunologically with whole cholera toxin or the individual A and B subunits. The enterotoxin caused increased cAMP activity (table V) in Chinese hamster ovary tissue culture cells and was capable of binding to GM1 ganglioside to the equivalent of 7 ng of cholera toxin (table VI).

The second, less active putative enterotoxin eluting with 0.55 M NaCl remains to be fully investigated.

Discussion

Following the emergence of detectable enterotoxin in the CFCF after incubation for 12 h at 37°C, with agitation at 100 oscillations/min in tryptone soya broth supplemented with yeast extract 0.6%, pH 7.5, the toxin became undetectable again just 5 h later. This interference with the functional activity of the toxin, probably the result of proteolytic breakdown, was accompanied by a change in the chromatographic profile of the CFCF; loss of enterotoxic activity corresponded with the appearance of a peak eluted with 0.22 M NaCl. Tests for enterotoxigenicity in routine isolates should, therefore, be carried out under strictly defined conditions of pH, temperature, agitation and time.

The infant mouse assay has been used in several studies^{5,6,9} to determine the enterotoxigenic activity of *Aeromonas* strains but reports of the factors essential for toxin production that may determine the outcome of these assays remain scarce. Our observations agree with those of Burke *et al.*⁶ and Turnbull *et al.*,⁹ who reported the importance of aeration of cultures during toxin preparation and the suitability of tryptone soya broth supplemented with yeast extract 0.6% as culture medium. We would also agree with Ljungh *et al.*¹² that toxin yields are higher after incubation at 37°C than at 20°C, but as they used different cultural conditions and demonstrated enterotoxic activity in the rabbit ileal loop test but not in the infant mouse assay, the comparison cannot be taken further. In the present study, β -haemolysin was produced rapidly by toxigenic strains during early logarithmic growth, but we confirmed that non-toxigenic strains⁹ remained inactive throughout prolonged incubation and under a range of cultural conditions.

Previous attempts to separate and purify aeromonas enterotoxin from culture filtrates by diverse methodologies have failed to identify a common protein.^{12–17} Poor yields and the instability of purified toxins have hindered their characterisation and the functional activities of single proteins have remained poorly defined.

Table VI. Binding characteristics of aeromonas enterotoxin to GM1 ganglioside

Preparation	Ganglioside GM1 binding assay		
	Absorbance (A ₄₁₀)	Estimated cholera toxin equivalent* (ng/ml)	Enterotoxigenicity in infant mouse (IW/BW ratio)
Crude aeromonas enterotoxin	0.18	5.8	0.0914
Crude aeromonas haemolysin	0.01	0.0	0.0677
Purified aeromonas enterotoxin	0.23	7.0	0.0956
Purified aeromonas haemolysin	0.02	0.0	0.062
Cholera toxin 50 ng	1.62	50.0	NT
Tryptone soya broth with 0.6% yeast extract	0.01	0.0	0.0529

NT, not tested.

Rabbit negative control sera gave no reaction.

* The estimated cholera equivalent values were estimated from a CT standard curve.

More than one functional activity, including enterotoxicity and haemolysis of erythrocytes have been assigned previously to single proteins¹³⁻¹⁵ but we found no evidence of this in the present study. The 55-kDa β -haemolysin described herein caused no fluid accumulation in the infant mouse assay and the existence of haemolytic activity in the CFCF did not correspond with either the emergence or decline of enterotoxic activity. In view of reports describing haemolytic enterotoxin,¹³⁻¹⁵ more than one type of aeromonas β -haemolysin may exist, and recent reports have indicated that several different antigenic types of haemolysin may occur.¹⁸

Purified aeromonas cytotoxic enterotoxin proved to be similar in size (15 kDa), to an active protein

described by Ljungh *et al.*¹² Its ability to stimulate cAMP production was demonstrated and, although it displayed some ability to bind to GM1 ganglioside, it did not cross-react immunologically with whole cholera toxin or cholera toxin subunits A and B. It is suggested, therefore, that aeromonas cytotoxic enterotoxin is a unique toxin which, in view of its molecular size, is unlikely to be divisible into subunits.

In addition to the 15 kDa enterotoxin a second enterotoxic fraction was eluted with 0.55 M NaCl. This fraction was also non-haemolytic and was immunologically distinct from the first enterotoxin and the haemolysin. Attempts to purify and characterise the second cytotoxic enterotoxin are in progress.

References

1. Lautrop H. *Aeromonas hydrophila* isolated from human faeces and its possible pathological significance. *Acta Pathol Microbiol Scand* 1961; **144**: 299-301.
2. Cumberbatch N, Gurwith MJ, Langston C, Sack RB, Brunton JL. Cytotoxic enterotoxin produced by *Aeromonas hydrophila*: relationship of toxigenic isolates to diarrhoeal disease. *Infect Immun* 1979; **23**: 829-837.
3. Gracey M, Burke V, Robinson J. *Aeromonas*-associated gastroenteritis. *Lancet* 1982; **2**: 1304-1306.
4. Agger WA, McCormick JD, Gurwith MJ. Clinical and microbiological features of *Aeromonas hydrophila*-associated diarrhoea. *J Clin Microbiol* 1985; **21**: 909-913.
5. Burke V, Gracey M, Robinson J, Peck D, Beaman J, Bundell C. The microbiology of childhood gastroenteritis: *Aeromonas* species and other infectious agents. *J Infect Dis* 1983; **148**: 68-74.
6. Burke V, Robinson J, Berry RJ, Gracey M. Detection of enterotoxins of *Aeromonas hydrophila* by a suckling mouse test. *J Med Microbiol* 1981; **14**: 401-408.
7. Ljungh A. *Aeromonas*-toxins and other virulence factors. Proceedings of the 1st International Workshop on *Aeromonas* and *Plesiomonas*, Manchester. 1986: 33-35.
8. Morgan DR, Johnson PC, DuPont HL, Satterwhite TK, Wood LV. Lack of correlation between known virulence properties of *Aeromonas hydrophila* and enteropathogenicity for humans. *Infect Immun* 1985; **50**: 62-65.
9. Turnbull PCB, Lee JV, Miliotis MD *et al.* Enterotoxin production in relation to taxonomic grouping and source of isolation of *Aeromonas* species. *J Clin Microbiol* 1984; **2**: 175-180.
10. Elliott RP, Clark DS, Lewis KH, Lundbeck H, Olson JC, Simonsen B (eds). Enumeration of mesophilic aerobes. In: *Micro-organisms in foods*. University of Toronto Press, ICMSF. 1978: 119-120.
11. Laemmli UK. Cleavage of structural proteins during the assembly of the head of the bacteriophage T4. *Nature* 1970; **227**: 680-685.
12. Ljungh A, Eneroth P, Wadstrom T. Cytotoxic enterotoxin from *Aeromonas hydrophila*. *Toxicon* 1982; **20**: 787-794.
13. Rose JM, Houston CW, Coppenhaver DH, Dixon JD, Kurosky A. Purification and chemical characterisation of a cholera toxin-cross-reactive cytolytic enterotoxin purified from a human isolate of *Aeromonas hydrophila*. *Infect Immun* 1989; **57**: 1165-1169.
14. Asao T, Kinoshita Y, Kosaki S, Uemura T, Sakaguchi G. Purification and some properties of *Aeromonas hydrophila* hemolysin. *Infect Immun* 1984; **46**: 122-127.
15. Potomski J, Burke V, Watson I, Gracey M. Purification of cytotoxic enterotoxin of *Aeromonas sobria* by use of monoclonal antibodies. *J Med Microbiol* 1987; **23**: 171-177.
16. Chakraborty T, Montenegro MA, Sanyal SC, Helmuth R, Bulling E, Timmis KN. Cloning of enterotoxin gene from *Aeromonas hydrophila* provides conclusive evidence of production of a cytotoxic enterotoxin. *Infect Immun* 1984; **46**: 435-441.
17. Potomski J, Burke V, Robinson J, Fumarola D, Miragliotta G. *Aeromonas* cytotoxic enterotoxin cross-reactive with cholera toxin. *J Med Microbiol* 1987; **23**: 179-186.
18. Chopra AK, Houston CW, Kurosky A. Genetic variation in related cytolytic toxins produced by different species of *Aeromonas*. *FEMS Microbiol Lett* 1991; **62**: 231-237.