A new multiplex PCR-based reverse line-blot hybridization (mPCR/RLB) assay for rapid staphylococcal cassette chromosome *mec* (SCC*mec*) typing

Lin Cai,^{1†} Fanrong Kong,^{2†} Qinning Wang,² Huiping Wang,³ Meng Xiao,⁴ Vitali Sintchenko^{2,5} and Gwendolyn L. Gilbert^{2,5}

Correspondence

Gwendolyn L. Gilbert
l.gilbert@usyd.edu.au

¹Department of Dermatology, Peking University People's Hospital, Beijing 100044, PR China

²Centre for Infectious Diseases and Microbiology, Institute of Clinical Pathology and Medical Research (ICPMR), Westmead, New South Wales, Australia

³Department of Dermatology, Tianjin Medical University General Hospital, Tianjin, PR China

⁴Life Science College, Peking University, Beijing, PR China

⁵Western Clinical School, University of Sydney, Sydney, Australia

The aim of this study was to develop a new discriminatory method for MRSA SCC*mec* typing based on multiplex PCR-based reverse line-blot hybridization (mPCR/RLB) assay to enable rapid identification and classification of MRSA SCC*mec* types in a clinical laboratory. Forty-five primer sets and 49 probes were designed and tested in uniplex PCR (uPCR) and mPCR/RLB. Probes were compared *in silico* to 14 whole-genome sequences and 18 partial SCC*mec* gene sequences of *Staphylococcus aureus* and complete genome and partial SCC*mec* genes of seven non-MRSA strains, including methicillin-susceptible *S. aureus* and methicillin-resistant coagulase-negative staphylococci. The method was tested on a set of 42 well-characterized reference MRSA strains. It identified all five epidemiologically relevant SCC*mec* types and 26 subtypes, including established and new subtypes of SCC*mec* III, IV (eight subtypes each) and V (three subtypes). The discriminatory power of mPCR/RLB SCC*mec* typing was similar to that of MLST and *spa* typing (Simpson indices of diversity of 0.916, 0.926 and 0.882, respectively; differences not statistically significant). The application of mPCR/RLB hybridization assay to MRSA SCC*mec* typing can improve the specificity, discriminatory power and throughput of the typing procedure. The detection of up to 43 mPCR products in a single hybridization assay transforms MRSA SCC*mec* typing from passive epidemiological library typing into a potential tool for near-real-time infection control surveillance and tracking of MRSA transmission in hospitals.

Received 12 November 2008
Accepted 5 May 2009

INTRODUCTION

Morbidity and mortality from methicillin-resistant *Staphylococcus aureus* (MRSA) infections and the emerging threat of community-associated MRSA highlight the importance of timely surveillance (Aires de Sousa & de Lencastre, 2004; Ma *et al.*, 2002). Methicillin resistance is conferred by the *mecA* gene, which is carried on a mobile genetic element known as the staphylococcal cassette chromosome *mec* (SCC*mec*) (Ito & Hiramatsu, 1998; Ito

et al., 1999). The combination of *mec* complex class and cassette chromosome recombinase (*ccr*) genes determines the SCC*mec* type (Ito *et al.*, 2004; Kondo *et al.*, 2007). Currently, seven SCC*mec* types (I–VII) and a number of subtypes have been described, ranging in size from 20.9 to 66.9 kb (Ito *et al.*, 2001, 2004; Daum *et al.*, 2002; Oliveira *et al.*, 2006a; Takano *et al.*, 2008). SCC*mec* types I and IV–VII are usually resistant only to β -lactam antibiotics, while types II and III are often resistant to multiple classes of antibiotics due to additional resistance genes integrated into SCC*mec* (Deurenberg & Stobberingh, 2008). MRSA SCC*mec* types I–III have been linked with healthcare-associated infections (Ito *et al.*, 2001) whereas community-acquired MRSA infections are typically caused by SCC*mec* types IV, V or VII (Ito *et al.*, 2004; Daum *et al.*, 2002; Takano *et al.*, 2008). Evidence has been emerging of

[†]These authors contributed equally to this work.

Abbreviations: mPCR/RLB, multiplex PCR-based reverse line-blot hybridization; MLST, multilocus sequence typing; MRSA, methicillin-resistant *S. aureus*; MSSA, methicillin-sensitive *S. aureus*; SCC*mec*, staphylococcal cassette chromosome *mec*; uPCR, uniplex PCR.

Table 1. Strains of 42 well-characterized meticillin-resistant *S. aureus* isolates used in the study

ID	Strain	<i>spa</i> type*	Clonal type†
1	FH53‡	t127	ST~1-SCCmec-I
2	COL§	t008	ST250-SCCmec-I
3	BK2464§	t002	ST5-SCCmec-II
4	E822485‡	t018	ST36-SCCmec-II
5	RPAH 18‡	t037	ST239-SCCmec-III
6	RPAH15‡	t037	ST239-SCCmec-III
7	HDG2§	t421	ST239-SCCmec-IIIB
8	14176-5710‡	t1959	ST~239-SCCmec-III
9	RDH81‡	t037	ST239-SCCmec-IIIA
10	AH1‡	t037	ST128-SCCmec-IIIA
11	AH13‡	t037	ST239-SCCmec-IIIA
12	HU25§	t138	ST239-SCCmec-IIIA
13	K704540‡	t037	ST~239-SCCmec-III
14	K711532‡	t037	ST~239-SCCmec-III
15	ANS46§	t037	ST239-SCCmec-III
16	PC8‡	t127	ST1-SCCmec-IV
17	FH43‡	t127	ST~1-SCCmec-IV
18	RPH 85‡	t127	ST~1-SCCmec-IV
19	RHH10‡	t127	ST~1-SCCmec-IV
20	B8-10‡	t711	ST~8-SCCmec-IV
21	PAH 58‡	t019	ST30-SCCmec-IV
22	E804531‡	t002	ST5-SCCmec-IV
23	CH97‡	T002	ST73-SCCmec-IV
24	RBH98‡	t202	ST93-SCCmec-IV
25	13792-4492‡	t202	ST~93-SCCmec-IV
26	IP01M1081‡	t216	ST59-SCCmec-IV
27	MW2§	t128	ST1-SCCmec-IV
28	CH16‡	t032	ST22-SCCmec-IV
29	CH69‡	t1963	ST~22-SCCmec-IV
30	DEN2988§	t008	ST8-SCCmec-IVA
31	F829549‡	t186	ST88-SCCmec-IV

potential associations between MRSA genotype and clinical outcomes (Lalani *et al.*, 2008).

SCCmec typing and subtyping provides an opportunity to detect and reduce the transmission of MRSA in healthcare settings (Ito *et al.*, 2004; Kondo *et al.*, 2007). Several multiplex PCR (mPCR)-based methods have been described to determine the structure of the SCCmec complex (Oliveira & de Lencastre, 2002; Zhang *et al.*, 2005; Kondo *et al.*, 2007; Milheirico *et al.*, 2007) but most require several reactions, produce long amplicons and are time consuming (Chongtrakool *et al.*, 2006; Kondo *et al.*, 2007), making them impractical for outbreak investigations or early-warning systems. Identification of the 46 major and minor SCCmec variants currently known relies on detection of the presence or absence of at least 22 of 34 binary targets (Stephens *et al.*, 2007). mPCR-based reverse line-blot hybridization assay (mPCR/RLB) can overcome the limitations of current methods by allowing rapid and simultaneous detection of a large number of different PCR products using membrane-bound sequence-specific probes (Kong & Gilbert, 2006). This technique has been

Table 1. cont.

ID	Strain	<i>spa</i> type*	Clonal type†
32	PAH 1‡	t019	ST30-SCCmec-IV
33	IP01M2046‡	t1958	ST78-SCCmec-IV
34	SJOG 30‡	t127	ST1-SCCmec-IV
35	RHH58‡	T127	ST~1-SCCmec-IV
36	IMVS 67‡	t008	ST8-SCCmec-V
37	RPH 74‡	t123	ST45-SCCmec-V
38	J710566‡	t065	ST45-SCCmec-V
39	B827549‡	t1784	ST~1-SCCmec-new
40	SN39‡	t127	ST~1-SCCmec-new
41	RPH2‡	t190	ST8-SCCmec-new
42	C801535‡	t325	ST88-SCCmec-new

**spa* type described by Cai *et al.* (2007).

†ST~, single nucleotide polymorphism (SNP) type as described previously using the computer program Minimum SNPs to compare with existing MLST data (Huygens *et al.*, 2004, 2006). SCCmec, staphylococcal cassette chromosome. ST, multilocus sequence type (MLST). Information provided by strain donors.

‡Provided by Dr Philip Giffard, Cooperative Research Centre for Diagnostics, Queensland University of Technology, Brisbane, Australia, and Dr Graeme Nimmo, Queensland Health Pathology Services (QHPS), Princess Alexandra Hospital, Brisbane, Australia, and tested in previous studies (Huygens *et al.*, 2004, 2006; Stephens *et al.*, 2006; Coombs *et al.*, 2004).

§Seven isolates, which had been used in previous studies (Oliveira *et al.*, 2006b), were provided by Professor Herminia de Lencastre, Instituto de Tecnologia Química e Biológica, Universidade Nova de Lisboa, Oeiras, Portugal; *spa* types identified in this study were identical with those previously reported for these strains (Cai *et al.*, 2007).

successfully applied to molecular typing of *Streptococcus agalactiae* (Kong *et al.*, 2005), *Streptococcus pneumoniae* (Kong *et al.*, 2006) and *Staphylococcus aureus* (Cai *et al.*, 2007).

The aim of this study was to develop a new discriminatory method for MRSA SCCmec typing based on mPCR/RLB suitable for the rapid identification and classification of the MRSA isolates in a clinical laboratory.

METHODS

Bacterial isolates. Seven well-characterized reference strains [Cai *et al.*, 2007; kindly provided by Professor Herminia de Lencastre, Instituto de Tecnologia Química e Biológica, Universidade Nova de Lisboa (UNL), Oeiras, Portugal] and 35 previously characterized clinical strains (provided by Drs Graeme Nimmo, Queensland Health Pathology Services, Princess Alexandra Hospital, Brisbane, Queensland, Australia, and Philip Giffard, Cooperative Research Centre for Diagnostics, Queensland University of Technology, Brisbane, Queensland, Australia) were used in this study (Table 1). Isolates were stored at -70°C , until DNA extracts were prepared, as previously described (Cai *et al.*, 2007); the latter were stored at -20°C until tested.

Primer and probe design. The mPCR primers (Kondo *et al.*, 2007) were modified to adjust T_m values to 60–65 °C and labelled with biotin. Immediate downstream or upstream sequences of the published sense and antisense primers were used to design antisense or sense mPCR/RLB probes as described previously (Kong & Gilbert, 2006). An inner primer was designed for each of the published sense and anti-sense primers with expected product size of around 100 bp. In total, 45 primer sets and 49 probes were assembled (some primer sets contained two probes). All primer sets were tested with each of the 42 reference strains using uniplex PCR (uPCR) and mPCR/RLB (Table 2).

uPCR and mPCR reactions. PCR amplifications were performed in a thermal cycler with HotStar Taq DNA polymerase (Qiagen). A 25 µl uPCR mixture was prepared follows: 2.5 µl 10 × PCR buffer with a final MgCl₂ concentration of 1.5 mM, 0.2 mM deoxynucleoside triphosphates, 12.5 µM of each of the forward and reverse primers, 2 µl template DNA, 0.2 µl Qiagen Hotstar Taq polymerase (5 U µl⁻¹) and molecular biology grade H₂O (Eppendorf) was added to a total volume of 25 µl. Two mPCR mixes were prepared based on uPCR results for the 42 reference strains: the first contained 39 primer pairs (primer sets 1–18, 21–24 and 29–45, Table 2), which produced positive uPCR results for at least one strain tested, and the second contained six primer pairs (primer sets 19, 20, and 25–28, Table 2) that were uPCR negative for all the strains tested, but showed *in silico* matches with one or more SCCmec sequences in GenBank.

Each mPCR mix comprised 2 µl DNA template and 3 µl 10 × PCR buffer with a final concentration of: 1.5 mM MgCl₂ (Qiagen), 0.2 mM dNTP, 1.5 U HotStar Taq DNA polymerase and 12.5 µM of each of the forward and reverse primers. Molecular biology grade H₂O (Eppendorf) was added to a total volume of 30 µl. PCR was performed according to the Qiagen HotStar Taq polymerase kit instructions: 95 °C for 15 min; 35 cycles of 94 °C for 30 s, 55 °C for 30 s, 72 °C for 45 s, and 72 °C for 10 min. PCR products were stained with SYBR safe DNA gel stain and visualized in a 2 % agarose gel.

Reverse line blot (RLB) hybridization. The RLB hybridization assay was performed as described previously (Kong & Gilbert, 2006; Cai *et al.*, 2007). Briefly, all probes, synthesized with a 5' amine modification at concentrations of 5–10 µM in 0.5 M NaHCO₃ (pH 8.4), were covalently linked to a Biotinylated C hydrophilic nylon 6,6 (0.45 µm) membrane (PALL Gelman Laboratory), which was negatively activated by incubation in freshly prepared 16 % (w/v) 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDAC) (Sigma). The membrane was placed in a miniblotted (MN45, Immunetics) on a support cushion. A volume of 20 µl of PCR products was added to 150 µl 2 × SSPE (Amresco) in 0.1 % SDS (Sigma), denatured at 100 °C for 10 min and cooled immediately on ice. Hybridization was conducted by adding 150 µl of the diluted denatured PCR products to alternate slots on the miniblotted, which was then incubated at 60 °C for 60 min. The membrane was washed twice in 2 × SSPE buffer in 0.5 % SDS at 60 °C for 10 min, incubated in peroxidase-labelled streptavidin conjugate (Roche) at 42 °C for 60 min, then washed twice in 2 × SSPE buffer in 0.5 % SDS at 42 °C for 10 min and twice in 2 × SSPE buffer at room temperature for 5 min. For detection, the membrane was incubated in 15 ml chemiluminescence blotting substrate ECL detection reagent (Amersham, GE Healthcare) for 2 min, covered with transparent plastic sheet in a film cartridge and exposed with chemiluminescence film (Amersham Hyperfilm ECL, GE Healthcare) for 4 min. RLB results were regarded as positive when a hybridization dot signal was clearly visible.

Quality control of mPCR/RLB results. To ensure reproducibility of mPCR/RLB results, positive and negative controls were run on each

membrane. The positive control was a mixture of DNA extracts from MRSA strains COL, BK2464, E822485, ANS46, DEN2988, PAH1, IP01M2046, RHH58, RPH74, B827549 and RPH2, which together contain all 43 probe targets; the negative control was master mix only, without DNA. Any signal produced by a test isolate that was similar to or stronger than the corresponding signal of the positive control was identified as positive. A signal produced by the negative control was assumed to be due to contamination and the assay was repeated. In practice, rare equivocal results would be resolved by uPCR for the relevant target, with sequencing of the amplicon, if required, as 'gold standard', but for the purpose of this study, all mPCR/RLB results were confirmed with uPCR.

SCCmec gene and staphylococcal whole-genome sequence analysis. Thirty-five SCCmec gene sequences published in NCBI GenBank, including 20 known SCCmec type (I–VI) and subtype sequences and 14 *Staphylococcus aureus* genome and 1 methicillin-resistant *S. epidermidis* genome sequences were analysed as reference sequences and tested *in silico* using the mPCR/RLB assay. These sequences (with strain numbers; and DDBJ/EMBL/GenBank accession numbers) were as follows: (a) 18 SCCmec sequences: type I.1 SCCmec (NCTC10442; AB033763); type II.1 SCCmec (N315; D86934); type II.2 SCCmec (JCSC3063; AB127982); type II.B SCCmec (AR05; AJ810123); type II.3 (type IIE) SCCmec (AR13.1/3330.2; AJ810120); type II.4 SCCmec (RN7170; AB261975); type III.1 SCCmec (HU25; AF422696); type III.1 SCCmec and SCCmercury (85/2082; AB037671); type IV.1 SCCmec (CA05 [JCSC1968]; AB063172); type IV.2 SCCmec (JCSC1978 [8/6-3P] AB063173); type IV.3 SCCmec (81/108 [MR108]; AB096217); type IV.C SCCmec (2314; AY271717); type IV.4 SCCmec (JCSC4469; AB097677); type IV.E SCCmec (AR43/3330.1; AJ810121); type IV.g SCCmec (M03-68; DQ106887); type IV.h (PL72; AF411936); type V SCCmec (JCSC3624 [WIS]; AB121219); type VI SCCmec (HDE288; AF411935); and (b) 14 *S. aureus* genomes: COL (CP000046), JH1 (CP000736), JH9 (CP000703), MRSA252 (BX571856), MW2 (BA000033), Mu3 (AP009324), Mu50 (BA000017), N315 (BA000018), USA300 (CP000255), USA300_TCH1516 (CP000730), RF122 (AJ938182), MSSA476 (BX571857), NCTC8325 (CP000253), Newman (AP009351).

In addition, the mPCR/RLB system was tested *in silico* against whole-genome and partial SCCmec sequences of seven non-MRSA strains: five methicillin-susceptible *S. aureus* (MSSA) [AJ938182, BX571857, CP000253, AP009351, AM983545 (SSCmec type II)], plus one methicillin-resistant coagulase-negative *Staphylococcus* [*S. epidermidis* CP000029 (SCCmec type II)] and *Staphylococcus pseudintermedius* [AM904732 (SCCmec type III)].

GenBank SCCmec sequence searches and alignments and design of primers and probes were conducted using BioManager (Sydney Bioinformatics, <https://www.angis.org.au/>) and Sigma (<http://www.sigma-genosys.com/calc/DNACalc.asp>) browsers. Results of BLASTN alignments of designed probe sequences with all currently available SCCmec types (I–VI) and subtype sequences (35 GenBank reference sequences) including whole-genome and partial SCCmec sequences of non-MRSA staphylococci were applied for *in silico* prediction of RLB hybridization patterns. The uPCR and mPCR/RLB results were recorded as binary data, exported into BioNumerics software (version 4.6, Applied Maths). Dendrograms were generated using the categorical coefficient and clustering by UPGMA algorithm.

Statistical analysis. The discriminatory power was estimated by calculating Simpson's index of diversity (Hunter & Gaston, 1988) and the 95 % confidence intervals (CI) were determined as described previously (Grundmann *et al.*, 2001).

Table 2. Primers and probes used in mPCR/RLB for SCC*mec* genotyping

Primer/ probe set	Primers (b) and probes (p) for mPCR/RLB*	GenBank accession no.	Gene or ORF	Reference SCC <i>mec</i> or SCC sequence(s)	Probe location	Primer and probe sequence (5'-3') [†]
1	mA1b mecAAp mecASp-1 mA2b	D86934 D86935 D86936 D86934	<i>mecA</i> <i>mecA</i> <i>mecA</i> <i>mecA</i>	Type II.1	<i>mec</i> complex <i>mec</i> complex	45813 TGCTATCCACCCTCAAACAGG ₄₅₈₃₃ 45860 GGTGTGCTTACAAGTGCTAATAATTC ₄₅₈₃₅ 46053 GGTAAAGGTTGGCAAAAAGATA ₄₆₀₇₅ 46098 AACGTTGTAACCACCCCAAGA ₄₆₀₇₈
2	1b ccrA1Ap 1An	AB033763 AB033763 AB033763	<i>ccrA1</i> <i>ccrA1</i> <i>ccrA1</i>	Type I.1	<i>ccr</i> complex	24838 CAATCAAAACCTATATCATCAATCAGTACG ₂₄₈₆₇ 24890 TGGAAAGCTTTTCGAATTTGAT ₂₄₈₆₉ 24944 TCATCAATATAGGGGTACAACATGTT ₂₄₉₁₁
3	β c1Sn ccrB1Sp β cb	AB033763 AB033763 AB033763	<i>ccrB1</i> <i>ccrB1</i> <i>ccrB1, ccrB2, ccrB3</i>	Type I.1, II.1, III.1	<i>ccr</i> complex	25421 CAGGCAAGTTAATGCTCCAG ₂₅₄₄₀ 25497 CGGACAACGTCAAAGAGC ₂₅₅₁₄ 25541 ATTGCCTTGATAATAGCCITCT ₂₅₅₁₈
4	2b ccrA2Ap 2An	D86934 D86934 D86934	<i>ccrA2</i> <i>ccrA2</i> <i>ccrA2</i>	Type II.1	<i>ccr</i> complex	26325 TAAAGGCATCAATGCACAAACACT ₂₆₃₄₈ 26378 TCTTTGCACACTTCTAATACTTTATCTTCA ₂₆₃₄₉ 26432 CGTTTGTCAATCGCACCT ₂₆₄₁₅
5	β c2Sn ccrB2Sp β cb	D86934 D86934 D86934	<i>ccrB2</i> <i>ccrB2</i> <i>ccrB1, ccrB2, ccrB3</i>	Type I.1, II.1, III.1	<i>ccr</i> complex	27168 TCGGAGTTTCTCAGAATTCTG ₂₇₁₈₇ 27226 ACGAGACGTGCCCCAAGAA ₂₇₂₄₃ 27261 ATTGCCTTGATAATAGCC ₂₇₂₄₄
6	A3b orfS04Ap	AB037671 AB037671	<i>ccrA3</i> <i>orfS04</i>	Type III.1	Adjacent to <i>ccr</i> complex	5486 AGCTCAAAAGCAAGCAATAGAAT ₅₅₀₈ 5533 TGAATATGTGCTTTGGAGCTA ₅₅₁₂
7	3An β c3Sn ccrA3Sp β cb	AB037671 AB037671 AB037671 AB037671	<i>orfS04</i> <i>ccrA3</i> <i>ccrA3</i> <i>ccrB1, ccrB2, ccrB3</i>	Type I.1, II.1, III.1	<i>ccr</i> complex	5583 TTCGATTATCTGTTCTGCCTGA ₅₅₆₂ 7173 TACAAATTCTAGCGAGTTTTTCAG ₇₁₉₆ 7241 ACCCGACGCGCTCAAGAA ₇₂₅₈ 7276 ATTGCCTTGATAATAGCC ₇₂₅₉
8	4.2b ccrA4Ap 4.2An	AF411935 AF411935 AF411935	<i>ccrA4</i> <i>ccrA4</i> <i>ccrA4</i>	Type VI	<i>ccr</i> complex	8741 AAAGGTATCAATGCACCAGAACTT ₈₇₆₄ 8780 ACAAGTGTGCGAGCACTTTACTTTC ₈₇₆₅ 8843 TGAGTATTGTGTCAATCAATCGATTA ₈₈₁₈
9	β 4.2Sn ccrB4Sp β 4.2b	AF411935 AF411935 AF411935	<i>ccrB4</i> <i>ccrB4</i> <i>ccrB4</i>	Type VI	<i>ccr</i> complex	9939 TGAGGAACCCATTATTGCTG ₉₉₅₈ 9985 AAAGCTTTATGGGATAAAGTACAGTT ₁₀₀₁₀ 10031 TTGCGACTCTCTTGGCGTTT ₁₀₀₁₂
10	Rb ccrCSp Fasn	AB121219 AB121219 AB121219	<i>ccrC</i> <i>ccrC</i> <i>ccrC</i>	SCC <i>mercury</i> , type V	<i>ccr</i> complex	16838 GCCTTTATAGACTGGATTATTCAAAAATA ₁₆₈₁₂ 16792 TCAGTGTTTGGCGTGAAAT ₁₆₈₁₀ 16741 ACGACAATTCTAAATCAAAAAGGA ₁₆₇₆₄
11	Ran ccrCAp Fb	AB121219 AB121219 AB121219	<i>ccrC</i> <i>ccrC</i> <i>ccrC</i>	SCC <i>mercury</i> , type V	<i>ccr</i> complex	16430 TTGAATTCATGAATAATATTAACACGT ₁₆₄₀₃ 16375 ATTTCCAACCTAATACCATTTTCGATT ₁₆₃₅₀ 16318 AAACGTCTATTACAAGATGTTAAGGATAAT ₁₆₃₄₇
12	mI6b mecR1Ap mA7An	D86934 D86934 D86934	<i>mecI</i> <i>mecR1</i> <i>mecR1</i>	Type II.1	<i>mec</i> complex	42866 CATAACTTCCCATTCTGCAGATG ₄₂₈₈₈ 42924 GAGTTAATATAATGGATAATAAACGTATGAAAT ₄₂₈₉₁ 42971 ATGGCGAAAAAGCACAAAC ₄₂₉₅₄
13	IS7b	AB033763	IS1272	Type I.1		28624 ATGCTTAATGATAGCATCCGAATG ₂₈₆₄₇

Table 2. cont.

Primer/ probe set	Primers (b) and probes (p) for mPCR/RLB*	GenBank accession no.	Gene or ORF	Reference SCC <i>mec</i> or SCC sequence(s)	Probe location	Primer and probe sequence (5'-3')†	
14	orfS034Ap	AB033763	<i>orfS034</i>	Type I.1, II.1, V	Adjacent to <i>mec</i> complex	28670TTTTGATTTTGTGATAGCCACA ₂₈₆₄₉	
	mA7An-1	AB033763	<i>orfS035</i>		<i>mec</i> complex	28726ATAAAGCATCTACTTTTGGATTTACTCT ₂₈₆₉₉	
	IS7Sn	AB033763	<i>mecR1</i>			31288CACCTTCTACACCTCCATATCA ₃₁₃₀₉	
	<i>mecASp</i> -2	AB033763	<i>mecA</i>			31395AAAGATAAAAAATTGTTCCACTTATTTTAATAG ₃₁₄₂₆	
15	mA7b	AB033763	<i>mecA</i>	Type I.1, II.1, V	Adjacent to <i>mec</i> complex	31450ATATACCAAACCCGACAACCTACA ₃₁₄₂₈	
	IS2(iS-2)Sn	AB121219	IS431			8670ACAGATCAGGCACCTTCAAC ₈₆₈₉	
16	IS431Sp	AB121219	IS431	Type V	J1	8721TTTAAACTTAAACCTGACTGTCTATTGT ₈₇₄₇	
	IS2(iS-2)b	AB121219	IS431			8772TGAGGTTATTCAGATATTTTCGATGT ₈₇₄₈	
	1a3b	AB033763	E007			Type I.1	5277TTTTAGGAGGTAATCTCCTTGATG ₅₃₀₀
	E007Ap	AB033763	E007			J1	5328TTTATAACATTGTCCTCTTTACCTTCAG ₅₃₀₁
17	E007Sp	AB033763	E007	Type I.1	J1	5387AATAAGCATACATGGAGACGTTTT ₅₄₁₀	
	1a4b	AB033763	E007			5431TTTTGCGTTTGCATCTCTACC ₅₄₁₁	
	4alb	AB063172	CQ002			Type IV.1	4726TTTGAATGCCCTCCATGAATAAAAT ₄₇₅₀
	CQ002Ap	AB063172	CQ002			J1	4777CGGGTAATATGAGAATAAATAAAGAAG ₄₇₅₁
18	4a1An	AB063172	CQ002	Type IV.1	J1	4844GCGTATTATGATACTAAGCAAAGTATTAATAATAG ₄₈₁₁	
	4a3Sn	AB063172	CQ002			5079TTGAAATCCTGGGAATCTATCACT ₅₁₀₂	
	CQ002Sp	AB063172	CQ002			5133ACATGCTGTAGTCATTTTATACTGTTTG ₅₁₆₀	
	4a3b	AB063172	CQ002			Type IV.2, II.3	5183AGCATATAGAAAAGATAGAAGTTCGAAAGA ₅₁₆₁
19	4b3b	AB063173	M001	Type IV.2, II.3	J1	2457AACCAACAGTGGTTACAGCTT ₂₄₇₇	
	M001Ap	AB063173	M001			2509AAATTTTAAACAATTCTAATTCAAATAAGTTG ₂₄₇₈	
	4b3An	AB063173	M001			2562ATCAAATTTTTTAGCAGATGTTGG ₂₅₃₉	
	4b4Sn	AB063173	M001			3041ATAGTGAACGTACTGATAACAAATTATTAGAT ₃₀₇₂	
20	M001Sp	AB063173	M001	Type IV.2, II.3	J1	3131ATCATAAAATTACAGCTTACGATATACAAA ₃₁₆₀	
	4b4b	AB063173	M001			3182CGGATTTTAGACTCATCCCAT ₃₁₆₁	
	4c4b	AB096217	CR008			Type IV.3	8266AAGAAAAGGAAATCGATGTCATTATAA ₈₂₄₀
	CR008Ap	AB096217	CR008			J1	8055TCCTACTCCTATAAAGAGGTTACAGAATTATT ₈₀₂₄
21	CR008Sp	AB096217	CR008	Type IV.3	J1	8208CAGATAGTATGGTACATATCCTTTAGGATT ₈₂₃₇	
	4c5b	AB096217	CR008			Type IV.4	7998CTGCATCCATTTCTCAGGAGTTAG ₈₀₂₁
	4d3b	AB097677	CD002			Type IV.4	2390AATTCACCCGTACCTGAGAA ₂₄₀₉
	CD002Ap	AB097677	CD002			J1	2440AGGATTTGGTGTATATAAAGGAGTGAAATA ₂₄₁₁
22	4d3An	AB097677	CD002	Type IV.4	J1	2496GCCTTTACACTAAGTACAGGGACATT ₂₄₇₁	
	4d4Sn	AB097677	CD002			3497TCTCTTAGAATTTTTTGTATATATTCATCTGA ₃₅₂₈	
	CD002Sp	AB097677	CD002			3575TGTACATCTCGTATGTTAGAAACATTATTTAAA ₃₆₀₇	
	4d4b	AB097677	CD002			3635GTCCAGAATGTGGTTATAAGATAGCTA ₃₆₀₉	
23	kdpB1b	D86934	<i>kdpB</i>	Type II.1	J1	12439TGAGATTACTTCAGAACCAGGTCAT ₁₂₄₁₅	
	kdpBAp	D86934	<i>kdpB</i>			12190GGCTATAGGGATTGCAGGG ₁₂₁₇₂	
	kdpBSp	D86934	<i>kdpB</i>			12388CAAACCAATCATTTTATCTAAAAATG ₁₂₄₁₃	

Table 2. cont.

Primer/ probe set	Primers (b) and probes (p) for mPCR/RLB*	GenBank accession no.	Gene or ORF	Reference SCCmec or SCC sequence(s)	Probe location	Primer and probe sequence (5'-3')†
25	kdpB2b 2b3b S01Ap	D86934 AB127982 AB127982	<i>kdpB</i> S01 S01	Type II.1 Type II.2	J1	12150 TAAACTGTGTACACGATCCAT ₁₂₁₇₁ 1497 GCTCTAAAAGTTGGATATGCG ₁₅₁₇ 1546 AAAGTAATCTATATGATCCCATTTTGA ₁₅₁₉
26	2b3An 2b4Sn S01Sp 2b4b	AB127982 AB127982 AB127982 AB127982	S01 S01 S01 S01	Type II.2	J1	1599 TTAGTATTTGTTACAATGCCCA ₁₅₇₆ 2906 ACAGTTAGAACATGTTAATAAAAAACATCG ₂₉₃₄ 2963 CAAATTAAGCAAACCTTACTGAATATTTAGC ₂₉₉₂ 3014 TGGATTGAATCGACTAGAATCG ₂₉₉₃
27	II4-3b RN06Ap II4-3An	AB261975 AB261975 AB261975	RN06 RN06 RN06	Type II.4	J1	11846 AAGTACCGCTGAATATTGATAGTGAT ₁₁₈₇₁ 11909 CCTATTAATTGAGATATTGTATGATAATATCCTT ₁₁₈₇₆ 11975 GGTTTTGTTTTTTCAATATAATACCTTG ₁₁₉₄₈
28	II4-1Sn RN06Sp II4-1b	AB261975 AB261975 AB261975	RN06 RN06 RN06	Type II.4	J1	13715 TTGAAATACATGTTTTTGAAGAATATATATTG ₁₃₇₄₆ 13797 AGAAATTGATTTAATAAGTTCCTCTGAAAATA ₁₃₈₂₈ 13850 TTTCTACTCTAATCCTAATCACCGAAC ₁₃₈₂₉
29	3a1b Z004Ap 3a1An	AB037671 AB037671 AB037671	Z004 Z004 Z004	Type III.1	J1	3333 ATGGCTTCAGCATCAATGAG ₃₃₅₂ 3381 AGTGCATTGCTTAATGATTAAATGTACT ₃₃₅₄ 3430 GACACCCGTATACAACGATGAC ₃₄₀₉
30	3a3Sn Z004Sp 3a2b	AB037671 AB037671 AB037671	Z004 Z004 Z004	Type III.1	J1	3734 ACTGCAGCGGCTTTAGGA ₃₇₅₁ 3793 GTTTAGATCATGGAATGGCACT ₃₈₁₄ 3835 ATATCCTTCAAGCGCGTTTC ₃₈₁₆
31	5a1b V024Ap 5a1An	AB121219 AB121219 AB121219	V024 V024 V024	Type V	J1	26564 ACCTACAGCCATTGCATTATG ₂₆₅₈₄ 26621 TCTTGATACATTTTAATATACTCACTAGTATTCGAT ₂₆₅₈₆ 26684 GCGAATATAAAATTCATATACTTTTCATCA ₂₆₆₅₅
32	5a2Sn V024Sp 5a2b	AB121219 AB121219 AB121219	V024 V024 V024	Type V	J1	27617 GGGATTTTCATCATCTTTTTAATGA ₂₇₆₄₀ 27671 AATTGATACTTATAAAATGTTAATAGCGCT ₂₇₇₀₀ 27722 TGTATACATTTTCGCCACTAGCT ₂₇₇₀₁
33	ermA1b ermAAp ermA1An	D86934 D86934 D86934	<i>ermA</i> <i>ermA</i> <i>ermA</i>	Type II.1	J2	35071 AATTTAGTGAACAATTTGTAACATTGA ₃₅₀₉₉ 35126 TATCGAAGGAACAATTTCTTTCTATT ₃₅₁₀₁ 35182 TAAAAACCAATTCGACAGG ₃₅₁₆₃
34	mN5Sn CN031Sp mN5b	D86934 D86934 D86934	CN031 CN031 CN031		J2	37724 CCTCCCATATATATAACAAGTTTAGG ₃₇₇₅₀ 37783 ATCCACTAAGTGTACTGAAAAATACCTG ₃₇₈₁₀ 37833 TTGCTTCGGGACTTACCTCTAGT ₃₇₈₁₁
35	cad4b cadBAp cad4An	AB037671 AB037671 AB037671	<i>cadB</i> <i>cadB</i> Z020-Z021 spacer	Type III.1	J2	16107 ATTGCGATTCTTTCCGATATGG ₁₆₁₂₈ 16156 CTATTTAATGCTACCAAAATAGTGGCTC ₁₆₁₂₉ 16213 TTCCGTAAATTGCACACATATAGTACTT ₁₆₁₈₆
36	mN5Sn Z023Sp mN5b	AB037671 AB037671 AB037671	Z023 Z023 CN030, CZ021	Type II.1, III.1	J2	17537 CCTCCCATATATATAACAAGTTTAGGA ₁₇₅₆₄ 17594 TGATCCACTAAGTGTACTGAAAAATACTTG ₁₇₆₂₃ 17646 TTGCTTCGGGACTTACCTCTAGT ₁₇₆₂₄
37	mA1b antAp	D86934 D86934	<i>mecA</i> <i>mecA</i>	Type II.1, III.1	J3	45813 TGCTATCCACCCTCAAACAGG ₄₅₈₃₃ 45859 GTGTGCTTACAAGTGCTAATAATTCA ₄₅₈₃₄

Table 2. cont.

Primer/ probe set	Primers (b) and probes (p) for mPCR/RLB*	GenBank accession no.	Gene or ORF	Reference SCC <i>mec</i> or SCC sequence(s)	Probe location	Primer and probe sequence (5′-3′)†		
38	ant1An mA1Sn antSp ant1b	D86934 D86934 D86934 D86934	<i>mecA</i> <i>ant(4)′</i> <i>ant(4)′</i> <i>ant(4)′</i>	Type II.1	J3	45914GGCATGAGTAACGAAGAATATAATAA ₄₅₈₈₉ 50668AGGCAAATGGCGTAATATTC ₅₀₆₈₇ 50724CTGTACAGGTAGCAATGGCA ₅₀₇₄₃ 50764GGTGCCATGTTGATTGGTCTG ₅₀₇₄₄		
39	mA1Sn-1 CZ036Sp pT181-2b	AB037671 AB037671 AB037671	CZ034-CZ036 spacer CZ034-CZ036 spacer CZ034-CZ036 spacer			Type III.1 Type I.1	J3	32761AATAATAAGGATAGCCATGGCTACAA ₃₂₇₈₆ 32815ATTTATAAATTCCAGAACTAGTTTACGATT ₃₂₈₄₄ 32869GGAGGTTTATTGTCACTACAATTGA ₃₂₈₄₇
40	la2b CE009Ap la1An	AB033763 AB033763 AB033763	<i>pls</i> CE009 CE009					Type I.1 SCC <i>mercury</i>
41	la2Sn plsSp la1b	AB033763 AB033763 AB033763	<i>pls</i> <i>pls</i> <i>pls</i>	J1	J1			
42	merA2b merAAp merA2An	AB037671 AB037671 AB037671	<i>merA</i> <i>merA</i> <i>merT</i>			SCC <i>mercury</i>	mer operon	
43	merGSn CZ046Sp merGb	AB037671 AB037671 AB037671	CZ046 CZ046 CZ046					SCC <i>mercury</i>
44	mN21b CZ055Ap	AB037671 AB037671	CZ055 CZ055	SCC <i>mercury</i>	J region in SCC <i>mercury</i>			
45	mN21An mN22Sn CZ056Sp	AB037671 AB037671 AB037671	CZ055 CZ056 CZ056			SCC <i>mercury</i>	J region in SCC <i>mercury</i>	
	mN22b	AB037671	CZ056					

*S, sense; A, antisense; b, biotin-labelled primers (at the 5' end); n, non-biotin-labelled primers; p, C⁶-amine-labelled-probes (at the 5' end). All primers labelled with biotin were based on previously published sequences (Kondo *et al.*, 2007). Non-biotin-labelled primers and probes were designed for this study.

†Numbers indicate sequence position in relevant GenBank sequences.

RESULTS AND DISCUSSION

SCCmec type characteristics of the reference strains in mPCR/RLB assays

All 42 reference strains were tested by uPCR and mPCR using all 45 primer sets (Table 2). The majority produced amplicons of the expected size in the uPCR, but six (primer sets 19, 20, 25–28) gave negative results. The two groups of primers (the 39 that generated and six that did not generate amplicon signals in the uPCR) were further studied in two separate mPCRs. Fig. 1 shows mPCR/RLB findings using the 39 primers that had given positive results in uPCR. As the other six primer sets did not produce signals with the 42 test isolates, they were removed from the membrane, to save space, and labelled onto a second membrane for use in subsequent studies.

The uPCR results were consistent with those of mPCR/RLB, with the following exceptions: primers $\beta c1Sn$ and βcb amplified SCCmec from strains FH53, SJOG30, RHH58, B827549 and SN39 in the uPCR but produced no signal in mPCR/RLB with probe ccrB1Sp (Fig. 1, lanes 1, 34, 35, 39 and 40, respectively), presumably due to a major sequence variation in the probe region. There were minor differences in intensities of signal dots but, since this is a binary typing system, clear signals similar to or stronger than positive controls on the membrane were regarded as positive. This approach was validated internally by the use of paired (sense/antisense) probes and by comparison of the signals within and between the probes. In our experience, a weak signal reflects minor sequence variation (1–2 bp) and an absent signal represents at least a 2 bp sequence change in the corresponding probe region, as suggested previously (Wang *et al.*, 2008). However, discordant signals between

Fig. 1. RLB patterns of 41 reference strains of MRSA. Lane P, artificial positive control, Lane N, negative control (PCR master mix). Lanes 1–42: 1, FH53; 2, COL; 3, BK2464; 4, E822485; 5, RPAH18; 6, RPAH15; 7, HDG2; 8, 14176-5710; 9, RDH81; 10, AH1; 11, AH13; 12, HU25; 13, K704540; 14, K711532; 15, ANS46; 16, PC8; 17, FH43; 18, RPH85; 19, RHH10; 20, B8-10; 21, PAH58; 22, E804531; 23, CH97; 24, RBH98; 26, IP01M1081; 27, MW2; 28, CH16; 29, CH69; 30, DEN2988; 31, F829549; 32, PAH1; 33, IP01M2046; 34, SJOG30; 35, RHH58; 36, IMVS67; 37, RPH74; 38, J710566; 39, B827549; 40, SN39; 41, RPH2; 42, C801535. [Note: strain 25 (13792-4492) was not included; it gave results identical to those of strain 24 (RBH98).]

probe pairs (one positive – weak or strong – and one negative) can occur but are rare. uPCR and sometimes sequencing, using the same primers as used for that target in the mPCR, can be used to determine whether the discrepancy is due to a modified target (amplicon present) or a non-specific cross-reaction (amplicon absent).

***In silico* comparison of probe sequences with SCCmec gene sequences**

The potential sensitivity of the probes designed in the study was assessed in an *in silico* experiment. The GenBank sequences of 14 *S. aureus* whole-genome sequences, 14 partial SCCmec sequences and seven non-MRSA genome and SCCmec sequences were downloaded and compared with probe sequences from this study using BLASTN pairwise alignments. The results are expressed as virtual RLB profiles in Fig. 2, in which the six probes that gave negative results for the 42 reference strains were also included. This comparative analysis demonstrated that each of the 45 probes found homologous matches with at least one GenBank MRSA SCCmec sequence. It suggested that the six additional probes/primer sets could be potentially useful for SCCmec typing and should be retained in the mPCR/RLB.

It should be noted that sequences of MRSA SCCmec type II 2 (lane 3), type IIB (lane 4), type II.4 (lane 6, accession no. AB261975), type III.1 (lane 7), type IV.4 (lane 13, accession no. AB097677), and type VI (lane 18, accession no. AF411935) were deposited in GenBank as partial-length sequences and may represent only the subtype-definitive regions of the SCCmec gene (Kondo *et al.*, 2007). The SCCmec type patterns identified by RLB in our 42 reference strains were generally consistent with those extrapolated from MRSA SCCmec gene sequences in the public domain. For example, the majority of SCCmec type IV strains (Fig. 1, lanes 15–26) had SCCmec profiles indistinguishable from that of the genome sequence ID 27 (Fig. 2, lane 27, CP000255). Altogether 15 of 42 strains (35%) had the same RLB patterns as those of corresponding GenBank sequences. However, there was some heterogeneity in the RLB patterns of other reference strains compared with GenBank sequences of the same SCCmec types. For example, RLB patterns of reference strains of SCCmec type III differed from the published MRSA SCCmec type III subtype 1 sequence (Fig. 2, lane 8, AB037671). This heterogeneity suggests the existence of distinct SCCmec type III subtypes (Fig. 2).

Theoretical RLB profiles of seven non-MRSA genome sequences are shown in Fig. 2. Four of five MSSA demonstrated no matches between probes and genomes, but one MSSA, and both methicillin-resistant coagulase-negative *Staphylococcus* strains, contained some SCCmec elements, as shown by matching of variable numbers of probes with their genomic sequences. These results are consistent with previous findings of SCCmec elements in MSSA and coagulase-negative staphylococci (Mombach

Pinheiro Machado *et al.*, 2007; Hanssen & Sollid, 2007). It has been postulated that SCCmec elements can be transferred from coagulase-negative staphylococci to *S. aureus* (Shore *et al.*, 2008). Our results suggest that our mPCR/RLB typing system could also be extended to study of the distribution of SCCmec elements in other staphylococci.

Discrimination of SCCmec genotypes by mPCR/RLB

The mPCR/RLB SCCmec typing system developed in this study was based on an mPCR typing scheme described by Kondo *et al.* (2007) using *ccr* and *mec* gene complexes types with further discrimination based on differences in junkyard region 1 and the presence or absence of integrated transposons and plasmids. Another recently developed SCCmec typing system used minimal subsets of targets on *ccr* and *mec* gene complexes and three junkyard regions to provide maximal discriminatory capacity (Stephens *et al.*, 2007). The rationales for these two strategies are different: the former is based on known SCCmec variants, whereas the latter maximizes discrimination, by recognizing single nucleotide polymorphisms (SNPs) and is not limited to known SCCmec types. Our system was designed as an extension of the Kondo scheme, to discriminate major known SCCmec types and subtypes rapidly. However, it can be easily modified to accommodate probes for newly identified SCCmec elements.

The set of 42 well-characterized MRSA reference strains included isolates representing 5 SCCmec types, 9 clonal clusters, 22 *spa* types, 7 *nuc* types, 9 *femA* types, 14 toxin types and 15 MLST types (Cai *et al.*, 2007). The mPCR/RLB assay classified reference strains into 26 SCCmec subtypes, including one of SCCmec type I (1 strain), two of SCCmec II (2 strains), eight of SCCmec III (11 strains), eight of SCCmec IV (21 strains) and three of SCCmec V (3 strains). MRSA SCCmec type IV and its variants have increased in importance in recent years because they have been identified in virulent community-acquired strains (Deurenberg & Stobberingh, 2008). Four of the previously unclassified strains demonstrated unique RLB patterns resembling SCCmec types IV and V (Fig. 3).

Different sequence-based typing methods targeting different types of gene may not be directly comparable but they are often proposed as alternatives to pulsed-field gel electrophoresis for strain typing. The mPCR/RLB SCCmec typing method showed similar discriminatory power to that of MLST and *spa* typing in this small but diverse strain collection, as indicated by Simpson indices of diversity of 0.916 (CI 88.1–95.1%) for mPCR/RLB typing, 0.882 (CI 85.2–91.2%) for MLST and 0.926 (CI 90.3–94.9%) for *spa* typing, respectively. Further investigation is required to determine its practical utility for SCCmec typing of a larger collection of clinical isolates. However, combined with GenBank SCCmec sequence data, this typing scheme also has the potential to identify novel

Fig. 3. Relationships between 42 MRSA strains determined by mPCR/RLB and other typing methods. The MLST and *spa* typing results were reported by Cai *et al.* (2007).

SCCmec types/subtypes among clinical MRSA isolates with increased accuracy and study relationships between them. Reference strain FH53 (lane 1, Fig. 2) was previously

reported to be SCCmec type I (Cai *et al.*, 2007). However, it appears to be similar to SCCmec IV reference and clinical strains as well as to the published SCCmec type IV sequence

(Fig. 3). There was also one genetic marker difference (probe β c2Sp) for strain MW2 between the mPCR/RLB assay and *in silico* results which could be attributed to the polymorphism of the binding site.

Advantages of the mPCR/RLB approach

The SCCmec gene typing initially proposed by Kondo *et al.* (2007) involves several mPCRs that produce amplicons of significant size (up to 15 kb) and requires gel-based detection. These features reduce the relevance of the typing scheme for infection-control-based interventions and tracking of introduction or transmission of different SCCmec types/subtypes. In contrast, the application of mPCR/RLB to the SCCmec typing described here enables the simultaneous screening of up to 43 samples using two mPCRs with a turnaround time (including culturing of isolates, DNA extraction, mPCR setup and running, and RLB hybridization) of about two working days. The preparation of the RLB membrane takes less than 2 h, and it can be reused at least 20 times based on our experience. The consumables cost is ~AU\$8 per isolate. The results of mPCR/RLB can be represented as binary data, which are easily transferred and shared between diagnostic and public health laboratories, making the SCCmec typing easier to harmonize and suitable for rapid epidemiological studies of large numbers of isolates.

In conclusion, this SCCmec typing system improves our capacity to monitor molecular evolution and spread of MRSA and contributes to the development of effective strategies to control its dissemination. The application of the mPCR/RLB hybridization assay to MRSA SCCmec typing enhances the specificity, discriminatory power and throughput of the typing procedure. The sequence detection of up to 43 mPCR products in a single hybridization assay can transform MRSA SCCmec typing from epidemiological library typing into a tool for near-real-time infection control surveillance and MRSA hospital transmission tracking.

ACKNOWLEDGEMENTS

The authors thank Professor Herminia de Lencastre and Drs Philip Giffard and Graeme Nimmo for kindly sharing MRSA isolates for the study.

REFERENCES

- Aires de Sousa, M. & de Lencastre, H. (2004). Bridges from hospitals to the laboratory: genetic portraits of methicillin-resistant *Staphylococcus aureus* clones. *FEMS Immunol Med Microbiol* **40**, 101–111.
- Cai, Y., Kong, F., Wang, Q., Tong, Z., Sintchenko, V., Zeng, X. & Gilbert, G. L. (2007). Comparison of single- and multilocus sequence typing and toxin gene profiling for characterization of methicillin-resistant *Staphylococcus aureus*. *J Clin Microbiol* **45**, 3302–3308.
- Chongtrakool, P., Ito, T., Ma, X. X., Kondo, Y., Trakulsomboon, S., Tiensasitorn, C., Jamklang, M., Chavalit, T., Song, J. H. & Hiramatsu, K.

- (2006). Staphylococcal cassette chromosome *mec* (SCCmec) typing of methicillin-resistant *Staphylococcus aureus* strains isolated in 11 Asian countries: a proposal for a new nomenclature for SCCmec elements. *Antimicrob Agents Chemother* **50**, 1001–1012.
- Coombs, G. W., Nimmo, G. R., Bell, J. M., Huygens, F., O'Brien, F. G., Malkowski, M. J., Pearson, J. C., Stephens, A. J. & Giffard, P. M. (2004). Genetic diversity among community methicillin-resistant *Staphylococcus aureus* strains causing outpatient infections in Australia. *J Clin Microbiol* **42**, 4735–4743.
- Daum, R. S., Ito, T., Hiramatsu, K., Hussain, F., Mongkolrattanothai, K., Jamklang, M. & Boyle-Vavra, S. (2002). A novel methicillin-resistance cassette in community-acquired methicillin-resistant *Staphylococcus aureus* isolates of diverse genetic backgrounds. *J Infect Dis* **186**, 1344–1347.
- Deurenberg, R. H. & Stobberingh, E. E. (2008). The evolution of *Staphylococcus aureus*. *Infect Genet Evol* **8**, 747–763.
- Grundmann, H., Hori, S. & Tanner, G. (2001). Determining confidence intervals when measuring genetic diversity and the discriminatory abilities of typing methods for microorganisms. *J Clin Microbiol* **39**, 4190–4192.
- Hanssen, A. M. & Sollid, J. U. (2007). Multiple staphylococcal cassette chromosomes and allelic variants of cassette chromosome recombinases in *Staphylococcus aureus* and coagulase-negative staphylococci from Norway. *Antimicrob Agents Chemother* **51**, 1671–1677.
- Hunter, P. R. & Gaston, M. A. (1988). Numerical index of the discriminatory ability of typing systems: an application of Simpson's index of diversity. *J Clin Microbiol* **26**, 2465–2466.
- Huygens, F., Stephens, A. J., Nimmo, G. R. & Giffard, P. M. (2004). *mecA* locus diversity in methicillin-resistant *Staphylococcus aureus* isolates in Brisbane, Australia, and the development of a novel diagnostic procedure for the Western Samoan phage pattern clone. *J Clin Microbiol* **42**, 1947–1955.
- Huygens, F., Inman-Bamber, J., Nimmo, G. R., Munckhof, W., Schooneveldt, J., Harrison, B., McMahon, J. A. & Giffard, P. M. (2006). *Staphylococcus aureus* genotyping using novel real-time PCR formats. *J Clin Microbiol* **44**, 3712–3719.
- Ito, T. & Hiramatsu, K. (1998). Acquisition of methicillin resistance and progression of multiantibiotic resistance in methicillin-resistant *Staphylococcus aureus*. *Yonsei Med J* **39**, 526–533.
- Ito, T., Katayama, Y. & Hiramatsu, K. (1999). Cloning and nucleotide sequence determination of the entire *mec* DNA of pre-methicillin-resistant *Staphylococcus aureus* N315. *Antimicrob Agents Chemother* **43**, 1449–1458.
- Ito, T., Katayama, Y., Asada, K., Mori, N., Tsutsumimoto, K., Tiensasitorn, C. & Hiramatsu, K. (2001). Structural comparison of three types of staphylococcal cassette chromosome *mec* integrated in the chromosome in methicillin-resistant *Staphylococcus aureus*. *Antimicrob Agents Chemother* **45**, 1323–1336.
- Ito, T., Ma, X. X., Takeuchi, F., Okuma, K., Yuzawa, H. & Hiramatsu, K. (2004). Novel type V staphylococcal cassette chromosome *mec* driven by a novel cassette chromosome recombinase, *ccrC*. *Antimicrob Agents Chemother* **48**, 2637–2651.
- Kondo, Y., Ito, T., Ma, X. X., Watanabe, S., Kreiswirth, B. N., Etienne, J. & Hiramatsu, K. (2007). Combination of multiplex PCRs for staphylococcal cassette chromosome *mec* type assignment: rapid identification system for *mec*, *cr*, and major differences in junkyard regions. *Antimicrob Agents Chemother* **51**, 264–274.
- Kong, F. & Gilbert, G. L. (2006). Multiplex PCR-based reverse line blot hybridization assay (mPCR/RLB) – a practical epidemiological and diagnostic tool. *Nat Protoc* **1**, 2668–2680.

- Kong, F., Ma, L. & Gilbert, G. L. (2005). Simultaneous detection and serotype identification of *Streptococcus agalactiae* using multiplex PCR and reverse line blot hybridization. *J Med Microbiol* **54**, 1133–1138.
- Kong, F., Brown, M., Sabananthan, A., Zeng, X. & Gilbert, G. L. (2006). Multiplex PCR-based reverse line blot hybridization assay to identify 23 *Streptococcus pneumoniae* polysaccharide vaccine serotypes. *J Clin Microbiol* **44**, 1887–1891.
- Lalani, T., Federsoiel, J. J., Boucher, H. W., Rude, T. H., Bae, I.-G., Rybak, M. J., Tonthat, G. T., Corey, G. R., Stryjewski, M. E. & other authors (2008). Associations between the genotypes of *Staphylococcus aureus* bloodstream isolates and clinical characteristics and outcomes of bacteremic patients. *J Clin Microbiol* **46**, 2890–2896.
- Ma, X. X., Ito, T., Tiensasitorn, C., Jamklang, M., Chongtrakool, P., Boyle-Vavra, S., Daum, R. S. & Hiramatsu, K. (2002). Novel type of staphylococcal cassette chromosome *mec* identified in community-acquired methicillin-resistant *Staphylococcus aureus* strains. *Antimicrob Agents Chemother* **46**, 1147–1152.
- Milheirico, C., Oliveira, D. C. & de Lencastre, H. (2007). Update to the multiplex PCR strategy for assigning of *mec* element types in *Staphylococcus aureus*. *Antimicrob Agents Chemother* **51**, 3374–3377.
- Mombach Pinheiro Machado, A. B., Reiter, K. C., Paiva, R. M. & Barth, A. L. (2007). Distribution of staphylococcal cassette chromosome *mec* (SCCmec) types I, II, III and IV in coagulase-negative staphylococci from patients attending a tertiary hospital in southern Brazil. *J Med Microbiol* **56**, 1328–1333.
- Oliveira, D. C. & de Lencastre, H. (2002). Multiplex PCR strategy for rapid identification of structural types and variants of the *mec* element in methicillin-resistant *Staphylococcus aureus*. *Antimicrob Agents Chemother* **46**, 2155–2161.
- Oliveira, D. C., Milheirico, C. & de Lencastre, H. (2006a). Redefining a structural variant of staphylococcal cassette chromosome *mec*, SCCmec type VI. *Antimicrob Agents Chemother* **50**, 3457–3459.
- Oliveira, D. C., Milheirico, C., Vinga, S. & de Lencastre, H. (2006b). Assessment of allelic variation in the *ccrAB* locus in methicillin-resistant *Staphylococcus aureus* clones. *J Antimicrob Chemother* **58**, 23–30.
- Shore, A. C., Rossney, A. S., O'Connell, B., Herra, C. M., Sullivan, D. J., Humphreys, H. & Coleman, D. C. (2008). Detection of staphylococcal cassette chromosome *mec*-associated DNA segments in multiresistant methicillin-susceptible *Staphylococcus aureus* (MSSA) and identification of *Staphylococcus epidermidis* *ccrAB4* in both methicillin-resistant *S. aureus* and MSSA. *Antimicrob Agents Chemother* **52**, 4407–4419.
- Stephens, A. J., Huygens, F., Inman-Bamber, J., Price, E. P., Nimmo, G. R., Schooneveldt, J., Munckhof, W. & Giffard, P. M. (2006). Methicillin-resistant *Staphylococcus aureus* genotyping using a small set of polymorphisms. *J Med Microbiol* **55**, 43–51.
- Stephens, A. J., Huygens, F. & Giffard, P. M. (2007). Systematic derivation of marker sets for staphylococcal cassette chromosome *mec* typing. *Antimicrob Agents Chemother* **51**, 2954–2964.
- Takano, T., Higuchi, W., Otsuka, T., Baranovich, T., Enany, S., Saito, K., Isobe, H., Dohmae, S., Ozaki, K. & other authors (2008). Novel characteristics of community-acquired methicillin-resistant *Staphylococcus aureus* strains belonging to multilocus sequence type 59 in Taiwan. *Antimicrob Agents Chemother* **52**, 837–845.
- Wang, Q., Kong, F., Jelfs, P. & Gilbert, G. L. (2008). Extended phage locus typing of *Salmonella enterica* serovar Typhimurium, using multiplex PCR-based reverse line blot hybridization. *J Med Microbiol* **57**, 827–838.
- Zhang, K., McClure, J. A., Elsayed, S., Louie, T. & Conly, J. M. (2005). Novel multiplex PCR assay for characterization and concomitant subtyping of staphylococcal cassette chromosome *mec* types I to V in methicillin-resistant *Staphylococcus aureus*. *J Clin Microbiol* **43**, 5026–5033.